ANTHOCYANINS EXTRACTION FROM PURPLE POTATO (GĂLĂNEȘTI BLUE-VIOLET VARIETY)

Florentina DAMȘA¹, Alexandru WOINAROSCHY², Carmen Liliana BĂDĂRĂU³

Varieties of pigmented potato (Solanum tuberosum L.) are an important source of anthocyanins. The aim of this work is to establish the optimal conditions for purple potato anthocyanins extraction. The anthocyanins were extracted from Gălăneşti blue-violet variety using the soaking in ethanol and 1% acidified ethanol. The highest yield (178.25mg/100g FW) of anthocyanins was reached at the temperature of 60° C, extraction time 60 min. in absolute ethanol.

Keywords: anthocyanins extraction, purple potato

1. Introduction

The interest for natural antioxidants, especially from fruits and vegetables, has increased in recent years. Epidemiological studies indicated that a high level of natural antioxidants (ascorbic acids, vitamin E, carotenoids and phenolics) in human everyday diet can protect against cardiovascular diseases, cataract, cancer and aging-related disorders [1].

Potato (*Solanum tuberosum L.*) is one of the principal food crops in the world and the tubers are a good source of carbohydrates (starch), proteins and vitamin C. As a product of plant origin they also contain secondary metabolites (phytochemicals) [2, 3, and 4]. Polyphenolic compounds are a large group of phytochemicals and depending on their chemical structure they can be divided into the following classes: flavonoids, phenolic acids, tanins, stilbenes and lignans [5]. Anthocyanins (classified as flavonoids) are responsible for the color found in the pigmented potatoes.

Considering that anthocyanins possess anti-oxidant, anti-microbial and interesting dye properties, these compounds are used in the food industry as a "natural" food additive. Several researchers have studied the anthocyanin contents

¹ Department of Chemical and Biochemical Engineering, University POLITEHNICA of Bucharest, National Institute of Research and Development for Potato and Sugar Beet, Brasov, Romania

² Department of Chemical and Biochemicalp Engineering, University POLITEHNICA of Bucharest, Romania, e-mail: a woinaroschy@chim.upb.ro

³ Department of Chemical and Biochemical Engineering, University POLITEHNICA of Bucharest, University Transilvania of Brasov, Faculty of Food and Tourism, Romania

of colored potato tubers. The pigments have been identified as anthocyanin derivatives [6, 7]. In fact, the first anthocyanin in a colored potato, malvidin-3-(pcoumaroryl-rutinoside)-5-glucose, was identified by Chmielewska (1936) still the year 1930. [8]

The aim of this study was to investigate the effect of temperature (30 - 60° C) extractions time (20 – 40 min) and solvent type (ethanol and 1% acidified ethanol) on anthocyanins extraction from purple potato (Gălănești blue-violet variety).



Fig. 1. Purple potato

2. Experimental

2.1. Anthocyanins extraction

Purple potato (Fig. 1) in amount of 5 g was homogenized in ethanol and 1% acidified ethanol (according with conditions presented in Table 1).

Table 1

(Galanesti blue-violet variety)							
Sample no.	Sample mass, g	Solvent	Time, min	Temperature, ℃			
P1	5	Ethanol	20	30			
P2	5	Ethanol	20	45			
Р3	5	Ethanol	20	60			
P4	5	Ethanol	40	30			
P5	5	Ethanol	40	45			

Conditions for anthocyanins extraction from Purple Potato

P6	5	Ethanol	40	60
P7	5	Ethanol	60	30
P8	5	Ethanol	60	45
P9	5	Ethanol	60	60
P10	5	1% acidified ethanol	60	30
P11	5	1% acidified ethanol	60	45
P12	5	1% acidified ethanol	60	60



Fig. 2 Extraction procedure applied for anthocyanins

Extracts were centrifuged (10 000 rpm, 15 min) and concentrated at 45°C. The extraction procedure of anthocyanin pigments from Purple Potato is schematically presented in Fig. 2.

2.2. Quantification of total anthocyanins

The total anthocyanins content were determined by the differential pH method based on the property of anthocyanin pigments to change the color with pH. Two dilutions of the same sample were prepared, the first one in potassium chloride buffer (0.025 M, pH 1.0), and the second one in sodium acetate buffer (0.4 M, pH 4.5), pH being adjusted with HCl. After equilibration at room temperature for 15 min, the absorbance of two dilutions was read at 510 nm and 700 nm. Total monomeric anthocyanins (mg cyanidin 3-glucoside equivalent/ 100 g Fresh Weight) were calculated as follows:

$$\% w/w = \frac{A}{\varepsilon L} MW DF \frac{V}{W_t} 100$$
 (1)

$$A = (A_{510nm} - A_{700nm})_{pH=1} - (A_{510nm} - A_{700nm})_{pH=4.5}$$
(2)

The significance of symbols used in these relations are:

%w/w – percentage weight/weight (grams of solute in 100 grams of solution) A – absorbance

 ϵ – molar extinction coefficient (26900 L/mol cm)

L – path length

MW – molecular weight (449.2 g/mol for cyanidin 3-glucoside)

DF - dilution factor

V – volume

Wt - sample weight

The total anthocyanin content (TAC) of purple potato extract is presented in Table 2.

Table 2

	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12
Т	47.1	68.7	81.1	44.41	55.10	115.43	86.0	139.5	178.25	59.08	73.32	150.
А	±	±	±	±	±	±	±	±	±	±	±	$73\pm$
C	2.43	3.02	2.72	2.95	3.61	3.42	3.15	3.80	4.23	3.24	3.61	2.68

Total anthocyanin content of purple potato extract

The influences of temperature, time and solvent on anthocyanins extraction are presented in Fig. 3-5.

Fig. 3 shows the influence of contact time between solvent and sample on anthocyanins extraction. The contact time has a positive influence on the content of total anthocyanin pigments (for a contact time of 20 min was obtained 81.1 ± 2.72 mg cyanidin 3-glucoside equivalent/ 100 g Fresh Weight and for 60 min was obtained 178.25 ± 4.23 mg cyanidin 3-glucoside equivalent/ 100 g Fresh Weight).



Fig. 3. Influence of contact time between solvent and sample on anthocyanins extraction

Fig. 4 shows the influence of temperature on anthocyanins extraction. The temperature has a positive influence on the total anthocyanins content (for a temperature of 30° C was obtained 86.0 ± 3.15 mg cyanidin 3-glucoside equivalent/ 100 g Fresh Weight and for 60° C was obtained 178.25 ± 4.23 mg cyanidin 3-glucoside equivalent/ 100 g Fresh Weight).



Fig. 4. Influence of temperature on anthocyanins extraction



Fig. 5. Influence of extraction solvent on the anthocyanin pigments

Fig. 5 shows the influence of extraction solvent on the anthocyanins pigments. Even if the extraction efficiency is higher in the case of ethanol (178.25 \pm 4.23 mg cyanidin 3-glucoside equivalent/ 100 g Fresh Weight) compared with 1% acidified ethanol (150.73 \pm 2.68 mg cyanidin 3-glucoside equivalent/ 100 g Fresh Weight), this is more stable in acidic medium.

3. Conclusions

The highest anthocyanin content was found at extraction temperature of 60°C, for an extraction time of 60 min. with ethanol as solvent. The lowest anthocyanin content was found at extraction temperature of 30°C, for an extraction time of 40 min. with ethanol as solvent. The anthocyanin content are lower when is used as solvent 1% acidified ethanol. The results are similar with literature reports about purple potato anthocyanin content determined through differential pH method. The disadvantage of this method is that it cannot provide information regarding the individual anthocyanins.

REFERENCES

- [1] Steffen L. M., Jacobs J. D. R., Stevens J., Shahar E., Carithers T., Folsom A. R., Associations of whole-grain, refined-grain. And fruit and vegetable consumption with risks of all-cause mortality and incident coronary artery disease and ischemic stroke: The Atherosclerosis in Communities (ARIC) Study, Am. J. Clin. Nutr., 78, 2003, p. 383-390
- [2] Augustin J., Variations in the nutritional composition of fresh potatoes, J. Food Sci., 40, 1975, p. 1295–1299
- [3] Lewis CE, Walker J. R. L., Lancaster J. E., Sutton K. H., Determination of anthocyanins, flavonoids and phenolic acids in potatoes. I: Colored cultivars of Solanum tuberosum L, J. Sci. Food Agric., 77, 1998, p. 45–57
- [4] Brown C. R., Antioxidants in potato, Am. J. Potato Res., 82, 2005, p. 163–172
- [5] Ignat I., Volf I., Popa V. I., A critical review of methods for characterization of polyphenolic compounds in fruits and vegetables, Food Chemistry, 126, 2011, p. 1821-1835
- [6] Rodriguez-Saona L. E., Giusti M. M., Wrolstad R. E., Anthocyanin pigment composition of red-fleshed potatoes, J. Food Sci., 63, 1998, p. 458–465
- [7] Eichhorn S., Winterhalter P., Anthocyanins from pigmented potato (Solanum tuberosum L.) varieties, Food Res. Intern., 38, 2005, p. 943–948
- [8] Chmielewska Z., Sur les colorants des pommes de terre violette 'Negresse', Bull. Soc. Chim., 5, 1936, p. 575–588.

es B: Chemistry and Materials Science **CIENTIFIC BULLETIN** iversity POLITEHNICA of Bucharest 2014 POLITEHNICA Volume 76 Issue 3 editura university in Romania. Its traditions are connected to the foundation of the first Along 190 years of existence, UPB represents one of the fundamental and prestigious institutions of the Romanian tertiary education, being the main source for the technical specialists of Romania. In the recent classification of the Romanian universities, our university is the first as far as the scientific appliquées de l'École Polytéchnique Roi Carol II. In accordance with the edittorial staff's claims, in this journal there have been published especially the results of the the teaching staff's original research. At that time, it different names according to the changes in the name of the institution : Bulletin de Mathématiques et de Physique pures et appliquées de l'École Organized on new bases, it primarily reflects the scientific activity of the University academic staff, but numerous papers have well-known authors University POLITEHNICA of Bucharest (UPB) is the most important technical The first issue our university journal was published in Bucharest on March 1929 under the name Bulletin de Mathématiques et de Physique pures et Our journal appeared with no interruption since 1929 till now under slightly 1954), Buletinul Institutului Politehnic București (1955 – 1992), University Polytéchnique (1929 – 1940), Buletinul Politehnicei din București (1941 | POLITEHNICA of Bucharest Scientific Bulletin (1992 – present). represented the third journal of this profile in the country. higher technical school by Gheorghe Lazar in 1818. Our bulletin is issued four times per year. production is concerned. from abroad.

MPRESS

ISSN 1454-2331

iversity POLITEHNICA of Bucharest

Scientific Bulletin www.scientificbulletin.upb.ro

> aterina ANDRONESCU litor-in-Chief

Corneliu BURILEANU Nicolae ŢĂPUŞ

Co-Editor

Mihaela Eleonora UNGUREANU, Sorin CIUCĂ Series Editors

Editorial Board

nstitute of Physical Chemistry, Aghia nstitute of Inorganic Chemistry, RWTH uropean Membrane Institute, Montpellier University POLITEHNICA of Bucharest Jniversity POLITEHNICA of Bucharest Jniversity POLITEHNICA of Bucharest University POLITEHNICA of Bucharest niversity POLITEHNICA of Bucharest niversity POLITEHNICA of Bucharest niversity POLITEHNICA of Bucharest iniversity of California, Los Angeles ihail Dumitru BARBOIU oannis G. ECONOMOU aula L. DIACONESCU Gheorghe MATEESCU niversité Montpellier II University of Angers Francois KAJZAR **Gheorghe MARIA** Alina Catrinel ION ina **BĂDĂNOIU** aniela BERGER Aachen University **III ENGLERT** ndreAYRAL larin BANE **Anton FICAI** Horia IOVU araskevi

Institute National Polytechnique de Déutsche Gesellschaft für Materialkunde, University POLITEHNICA of Bucharest Centre Interuniversitaire de Recherche et des Sciences University POLITEHNICA of Bucharest University POLITEHNICA of Bucharest University POLITEHNICA of Bucharest Université Joseph Fourier de Grenoble Ingénierie de Matériaux de Toulouse **Doru Michael STEFĂNESCU** George PAPANICOLAOU TNO Science and Industry Florin STEFĂNESCU nstitute National University of Alabama Wim H. SILLEKENS Jean Pierre MILLET Peter Paul SCHEEP **Eric SAINT-AMAN Bernard VIGUIER** Frankfurt am Main Appliquées de Lyon University of Patras Angela POPESCU **Maria NICOLAE** Francis MAURY **Teodor VISAN Foulouse**

Registered in

Case Western University, Cleveland, Ohio

Ulrich's International Periodicals Directory, Inspec, Scopus, Cambridge Scientific Abstracts, Chemical Abstracts Service - Division of American Chemical Society, Elsevier Sciences's Bibliographic Databases, Engineering Village, Compendex, Metadex, Engineered Materials Abstracts

Florentina Neli MINCĂ, Anamaria CENJA Sub-editors

ISSN 1454 - 2331

University POLITEHNICA of Bucharest

SCIENTIFIC BULLETIN

founded in 1929

SERIES B: Chemistry and Materials Science

2014 ssue 3 Volume 76

0
+
I
e
+
I
0

CHEMISTRY

3 11 33 43 55 67 15 80 66 113 123 21 Improvements in quality of histidine-modified enhanced green fluorescent protein that Mihai Rissdörfer, Katharigatta Narayanaswamy Venugopala, Susanta Kumar Nayak, Florin electrode processes in ionic liquid solvents as mixtures of choline chloride with urea, Improvement in the determination of traces of nitrate and nitrite in natural mineral Livia Alexandra Gugoaşă, Raluca-Ioana Stefan-Van Staden, Grzegorz Bazylak, Jacobus Iuliana Moldoveanu, Raluca-Ioana Stefan-Van Staden, Jacobus Frederick Van Staden, Florica Vaja (Dumitru), Cornelia Guran, Denisa Ficai, Anton Ficai, Ovidiu Oprea Physicochemical, crystallography and DFT calculations on biologically active Amperometric microsensors based on inulins for the assay of L-T3 and L-T4 Analysis of L-thyroxine and 3,3',5-triiodo-L-thyronine using potentiometric Anthocyanins extraction from purple potato (Gălănești blue-violet variety) Judit Petres, Emese-Éva Bálint, Mária Szabó, László Szilágyi, Beáta Ábrahám Cytotoxic effects of ZnO nanoparticles incorporated in mesoporous silica Extraction of vegetable oils from ground seeds by percolation techniques Spectrofluorimetric analysis of cefotaxime sodium by using 4-fluoro-7-Alexandrina Țane, Ileana Rău, Aurelia Meghea, Andreea Daniela Nistor Mihaela Nour, Daniela Dună, Mirela Ioniță, Viorel Feroiu, Dan Geană Morpho - structural control of functionalized DNA based systems Florentina Damşa, Alexandru Woinaroschy, Carmen Liliana Bădărău Mariana Lili Mareș Badea, Anca Cojocaru, Liana Anicăi Daniela Știrbeț, Ioana Vasilescu, Gabriel-Lucian Radu Integrated process for y-butyrolactone production binds copper ions with nanomolar affinity nitrobenzofurazan as derivatization agent Frederick Van Staden, Gabriel-Lucian Radu Ion Ion, Daniela Bogdan, Alina Catrinel Ion Ahtesham Javaid, Costin Sorin Bildea ethylene glycol or malonic acid waters by ion chromatography dihydropyrimidine analogues Gabriel Lucian Radu microsensors Dinu Badea

> This issue appears thanks to the careful supervision of prof. Eleonora Mihaela UNGUREANU and prof. Sorin CIUCĂ

ISSN 1454-2331