

THE NPK FERTILIZATION EFFECTS OF TUBERS STARCH, DRY MATTER AND REDUCING SUGAR CONTENT

Nina BĂRĂSCU¹, Maria IANOȘI¹, Marcel M. DUDA², Edward MUNTEAN²

¹National Institute of Research and Development for Potato and Sugar Beet Brașov, Street Fundăturii no 2, Brașov, Romania

²University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, Calea Mănăștur 3-5, Postal code 400372, Cluj-Napoca, Romania

Corresponding author email: nina.barascu@gmail.com

Abstract

The content of potato tubers in dry matter, starch and reducing sugar determines their quality characteristics. This study shows how the basic fertilization with different nitrogen doses and NPK ratios influence the content of potato tubers in dry matter, starch and reducing sugar.

The researches were conducted in the years 2013-2014 in the experimental field of INCDCSZ Brașov, located in Brașov Depression, on a cernoziomoid soil at an altitude of 520 m. For the two potato varieties used in the experiment, Christian and Roclas, was made a basic NPK fertilization with two different ratios 1: 1: 1 and 1: 0.9: 2 and two different doses of nitrogen, 100 and 200 kg/ha.

Fertilization ratio 1:0.9:2 as against 1:1:1 ratio had significant effect on reducing the starch content for both varieties, while the reducing of dry matter content is significant only for Christian variety. For both varieties, there were no significant differences in reducing sugar. The higher NPK ratio and nitrogen dose of 200 kg/ha determined reduction of starch in the two years, on average with 0.9-1.2%. Increasing the dose of nitrogen determined a significant decrease in the reducing sugar content only at Christian variety.

The decrease in starch content from tubers of Christian variety with increasing potassium ratio and nitrogen dose is more evident and statistically significant in 2014. For Roclas variety there is the same tendency, differences between variants in the two years are lower.

In the case of the dry matter content differences due to experimental years are stronger and are manifested mostly for dose N200 both at Christian as well as the Roclas. For the tubers reducing sugar content were found significant differences between the two years for both varieties. To all variants of fertilization the content level of reducing sugar was lower in 2013.

Key words: dry matter, fertilization, potato, starch, reducing sugar.

INTRODUCTION

Potato culture has a great potential of production, which can be achieved if ecological and technological conditions are ensured. Potato crop has strict requirement for a balanced fertilization management, without which the crop's growth and development of the crop are poor and both yield and quality of tubers are diminished (Imas and Bansal, 1999). Plant nutrition is a key instrument for managing potato quality (Achten, 2005). The content of reducing sugars, dry matter, and starch in potato tubers are important properties for potato processing (De la Fuente, 2011) because quality of final products and efficiency of processing depend on them (Hasse, 2004). Nitrogen is the dominant nutrient taken up by

potato and plant nitrogen requirements are greater than other nutrient (El-Galil, 2006). A sufficient amount of nitrogen is important for achieving quality objectives for processing potatoes (Zebarth, 2012). Tuber specific gravity decrease if more nitrogen is available than needed for growth (Westermann et al., 1994; Atkinson et al., 2003).

Natural or chemical fertilization and soil type on which potatoes are grown may influence the tubers sugar content. Plants adequately fertilized with nitrogen had tubers with lower reducing sugar concentration at harvest (Kumar et al., 2004).

Potatoes require large amounts of soil K, since this nutrient is crucial to metabolic functions such as movement of sugars from the leaves to

the tubers and the transformation of sugar into potato starch (Mikkelsen, 2006).

Phosphorus applications may improve specific gravity when soil test phosphorus levels are low (Laboski and Kelling, 2007).

A deficiency of phosphorus and potassium in the soil may lead to decreased synthesis of starch and sugar content increases (Mureşan, 1999). Potassium plays an important role in the starch structure and reduces the percentage of reducing sugars (Ianoşi, 2002).

Among the major nutrients, potassium not only improves yields but also affects some of the tuber quality parameters like percentage of dry matter and starch content (Imas and Bansal, 1999) and the fertilizer potassium source is known to affect tuber specific gravity (Westermann et al., 1994).

Potassium contributes to various aspects of tuber quality and the balance between nitrogen and potassium supply is of particular importance for potato crop (El-Latif, 2011).

The aim of this work was to assess the effects of different NPK fertilization on the tubers starch, dry matter and reducing sugar content.

MATERIALS AND METHODS

The experimental design was based on the knowledge acquired in the domain of fertilization, on a black earth soil from Braşov and according to current practices of fertilization in the region. In these polifactorial experience we propose levels of fertilization and different NPK ratios, for Roclas and Christian varieties created at N.I.R.D.P.S.B. Brasov.

The chosen levels of nitrogen fertilization, of 100 and 200 kg N/ha are situated at the lower and upper limit of OSPA Braşov recommendations valid for autumn potato crops. These doses are recommended for production of 20 t/ha and 40 t/ha under non-irrigated conditions.

Two complex fertilizers given before planting: C15-15-15, for NPK ratio of 1:1:1 and C5:10:22, supplemented with ammonium nitrate through for obtaining a 1:0.9:2 NPK ratio.

The experiment was done in Braşov in a non-irrigated crop, studied during 2013-2014. The

research was conducted in two years with very different growth conditions in terms of climate. In 2013 the vegetation period was warm, with an average temperature of 16°C and 422.2 mm precipitation, compared to the year 2014, at which the average air temperature was 15.3°C and there has been rainfall of 505.1 mm (Figures 1 and 2).

In 2014, the amount of rainfall during the growing season exceeded the multiannual average (MMA) and the average temperature during the growing season was close to the multiannual value ensuring high yields in both studied varieties.




Figure 1. Average temperatures – Braşov 2013-2014




Figure 2. Average rainfall – Braşov 2013-2014

After harvesting and before cold storage of potato tubers, we made determinations on experimental variants and repetitions, regarding the tuber starch content, using the Polikeit balance. Also we determined the dry matter content of tubers by drying them in an oven at 105° C for 4 hours. Tubers content, from experimental variants, in reducing sugars (glucose, fructose) was determined by the

chromatographic method using Shimadzu HPLC system.

RESULTS AND DISCUSSIONS

On average, for the two experimental years, varieties differed in starch content, dry matter and reducing sugar before storage (Table 1.). In Christian variety there were recorded on average lower levels of starch, dry matter and reducing sugar (15.0%, 23.9% and 0.18%) compared to the Roclas variety (16.5%, 24.8% and 0.34%). Standard deviations indicate moderate variation of values calculated for starch and dry matter and high variation for reducing sugar.

The fertilization ratio 1:0.9:2 compared with that of 1:1:1 had significant effect on reducing the starch content for both varieties, while the reducing of dry matter content was significant only for Christian variety. For both potato

varieties there were no significant differences in reducing sugar. The higher NPK ratio and nitrogen dose of 200 kg/ha determined reduction of starch in the two years, on average with 0.9-1.2%

Increasing the dose of fertilizer from N100 to N200 determined a decrease in the starch content for both varieties, with no significant differences for dry matter. Although with increasing nitrogen level it can be noted a decrease for the reducing sugar content; decrease was significant only for Christian variety.

The Duncan test revealed that for the lowest level of fertilization (N100 NPK 1:1:1 ratio) was recorded the highest starch content, 16.1% for Christian variety and 17.7% for Roclas variety, while for variant with the highest level of fertilization (N200 1:0.9:2 ratio), the starch content for the two varieties decreased to 13.9%, for Christian and 15.6% la Roclas.

Table 1. Mean effects of different with NPK ratios and dozes fertilization on tubers starch, dry matter and reducing sugar content from Christian and Roclas varieties (Braşov 2013-2014)

Ratio NPK	Doses N Kg/ha	Starch %		Dry matter %		Reducing sugar %	
		Christian	Roclas	Christian	Roclas	Christian	Roclas
1:1:1	100	16.1 a	17.7 a	24.6 a	25.5 a	0.21 ab	0.41 a
	200	14.9 ab	16.2 bc	24.5 a	24.8 a	0.10 b	0.28 a
1:0.9:2	100	15.2 ab	16.5 b	23.7 a	24.7 a	0.29 a	0.33 a
	200	13.9 b	15.6 c	23.0 a	24.2 a	0.13 b	0.32 a
Means							
Ratio 1:1:1		15.5	17.0	24.6	25.1	0.16	0.35
Ratio 1:0.9:2		14.6 ^o	16.1 ^o	23.3 ^o	24.4	0.21	0.33
N 100 kg/ha		15.7	17.1	24.1	25.1	0.25	0.37
N 200 kg/ha		14.4 ^o	15.9 ^o	23.8	24.5	0.12 ^o	0.30
Mean		15.0	16.5	23.9	24.8	0.18	0.34
Standard deviation		1.5	1.0	1.9	1.9	0.16	0.21
LDS(variants)5%		2.0%	1.0%	2.7%	2.8%	0.21%	0.31%
DL (ratio) 5%		1.1%	0.7%	1.3%	1.4%	0.11%	0.15%
DL (doses N) 5%		1.0%	0.6%	1.4%	1.4%	0.10%	0.15%

On average fertilization variants were not significantly differentiated for the varieties researched, as regards the dry matter content.

The average content of reducing sugar of variants studied presented significant differences only for Christian variety. To this variety, for both fertilization ratios passing from N100 at N200 strongly reduced the content of reducing sugar (from 0.21% to 0.10% for NPK 1:1:1 and from 0.29% to 0.13% for NPK 1:0.9:2).

The content of starch, dry matter and reducing sugar of tubers was significantly influenced by growing conditions.

Due to the more favorable climatic hydro and thermal conditions alongside with higher yields for Christian variety accumulation of starch in tubers was significantly lower (16.2% as against 13.9%) in 2014 compared with 2013 (Figure 3.). On average, for Roclas variety, was maintained the high starch content from tubers, there are no significant differences between the two years (16.8% and 16.2%).

The decrease in starch content from tubers of Christian variety with increasing potassium ratio and nitrogen dose is more evident and statistically significant in 2014. For Roclas variety there is the same tendency, but the

differences between variants in the two years are lower.

In the case of the dry matter content, the differences due to experimental years are stronger and are manifested mostly for the N200 dose, for both varieties (Figure 4).




Figure 3. Comparison of the average starch content of tubers for fertilization variants - Braşov, 2013-2014




Figure 4. Comparison of the average dry matter content of tubers for fertilization variants - Braşov, 2013-2014




Figure 5. Comparison of the average reducing sugar content of tubers for fertilization variants - Braşov, 2013-2014

For the two studied years, correlations were made between the content of starch, dry matter and reducing sugar from tubers and total

production from tubers. Pearson correlation coefficients for Christian and Roclas varieties are presented in Table 2.

Table 2. Pearson correlation coefficients for total yield, starch, dry matter and reducing sugar content of tubers (Braşov, 2013-2014)

	Christian			Roclas		
	Starch	Dry matter	Reducing sugar	Starch	Dry matter	Reducing sugar
Starch	-	0.772**	-0.246	-	0.677**	-0.167
Dry matter	0.772**	-	-0.421*	0.677**	-	-0.331
Reducing sugar	-0.246	-0.421*	-	-0.044	-0.218	-
Total yield	-0.638**	-0.454**	0.488**	-0.315	-0.369*	0.689**

For both varieties the correlation between content of starch and dry matter is positive and statistically assured ($r = 0.772^{**}$ for Christian and $r = 0.677^{**}$ for Roclas). Correlations between reducing sugar, starch and dry matter for both varieties are negative. Only for Christian variety correlation between dry matter and reducing sugar content was statistically assured ($r = -0.421^*$). Correlations of total production with concentration of starch from tubers was significant for Christian variety ($r = -0.638^{**}$). The sense of correlations indicates very strong decreases of starch content due production increases, especially in 2014.

The correlations between the total production with starch content and dry matter from tubers were negative and statistically assured, for Christian variety and for Roclas variety correlations were lower. The correlations of the total production with reducing sugar were for both varieties positive and statistically assured.

CONCLUSIONS

Very different climatic conditions in those two years determined high yield differences from one year to another, and the accumulation of starch and dry matter in tubers was significantly lower in 2014 comparing with 2013.

On average, in both varieties, the highest starch content, was accumulated by the variants with fertilization N100: P100: K100 and on those two years, the differences due fertilization variants were not provided statistical for dry matter content of tubers. For 2014, the fertilization variants with high nitrogen level led to significant decreases in dry matter

content for both fertilizing ratios, for both varieties.

Both potato varieties revealed a significantly higher content of reducing sugar in 2014 comparatively with 2013.

ACKNOWLEDGEMENTS

This paper was published under the frame of European Social Fund, Human Resources Development Operational Programme 2007-2013, project no. POSDRU/159/1.5/S/132765.

REFERENCES

- Achten Vincent T.J.M., 2005. Technology developments in potato yield and quality management. In: Havercort A.J., Struik P.C., Eds., *Potato in progress: science meets practice*. Wagening Academic Publishers The Netherlands.
- Atkinson D., Geary B., Stark J., Love S., Windes J., 2003. Potato varietal responses to nitrogen rate and timing., Idaho Potato Conference, January 22
- De La Fuente L., Varas J., Mendez S., Elita Lopez M.A., 2011. Characterization of the content of reducing sugars, total sugars and starch in potato varieties and clones grown in Osomo-Chile, International Congress on Engineering and Food „Food process Engineering in a Changing World”, Athens, Greece.
- El-Galil Ali Abd, 2006. Fertilizer-n dynamics in the soil and yield response of potatoes as affected by methods of N-application, *Journal of Applied Sciences Research*, 2(9): 613-623.
- El-Latif Abd K.M., Osman E.A.M., Abdullah R., Abd El Kader N., 2011. Response of potato plants to potassium fertilizer rates and soil moisture deficit, *Advances in Applied Science Research*, 2(2): 388-397
- Hasse Norbert U., 2003. Estimation of dry matter and starch concentration in potatoes by determination of under-water weight and near infrared spectroscopy, *Potato Research* 46 9(2003/4) 117-127.
- Ianoşi I.S., 2002. Bazele cultivării cartofului pentru consum, Ed. Phoenix, Braşov.

- Imas P., and Bansal S.K., 1999. Potassium and integrated nutrient management in potato, Global Conference on Potato, 6-11 December 1999, New Delhi, India.
- Kumar Dinesh, B.P. Singh and Parveen Kumar, 2004. An overview of the factors affecting sugar content of potatoes, *Annals of Applied Biology*, 145, p. 247-256.
- Laboski Carrie A.M., Lelling K., 2007. Influence of fertilizer management and soil fertility on tuber specific gravity: a review, *American Journal of Potato Research*, 84:283-290
- Mikkelsen R., 2006. Best management practices of profitable fertilization on potatoes, *Better Crops*, vol. 90, nr.2.
- Mureșan S., 1999, Calitatea cartofului pentru consum. Calitatea culinară a tuberculilor de cartof, *Cartoful în România*, vol. 9, nr.2, aprilie-iunie.
- Westermann D.T., James D.W., Tindall T.A., Hurst R.L., 1994. Nitrogen and potassium fertilization of potatoes: sugars and starch, *American Potato Journal*, vol. 71, issue 7, p. 433-453
- Zebarth B.J., Belanger G., Cambouris A.N., Ziadi N., 2012. Nitrogen fertilization strategies in relations to potato tuber yield, quality and crop N recovery, in *Sustainable Potato Production: Global Case Studies*, editors: Z. He, R. Larkin, W. Honeycutt, part IV, chapter 10, p. 165-186.

SCIENTIFIC PAPERS
SERIES A. AGRONOMY
VOLUME LIX, 2016

UNIVERSITY OF AGRONOMIC SCIENCES
AND VETERINARY MEDICINE OF BUCHAREST
FACULTY OF AGRICULTURE

SCIENTIFIC PAPERS
SERIES A. AGRONOMY

VOLUME LIX

2016
BUCHAREST

SCIENTIFIC COMMITTEE

- Costică CIONTU – Faculty of Agriculture, USAMV Bucharest, Romania
- Cristian HERA – Romanian Academy
- Iovu-Adrian BIRIȘ – Faculty of Agriculture, USAMV Bucharest, Romania
- Raimundo CABRERA – University of La Laguna, Phytopathology Unit, Spain
- Ionela DOBRIN – Faculty of Agriculture, USAMV Bucharest, Romania
- Marin DUMBRAVĂ – Faculty of Agriculture, USAMV Bucharest, Romania
- Mihail DUMITRU – ICPA Bucharest, Romania
- André FALISSE – University of Liège, Gembloux Agro-Bio Tech, Belgium
- Beatrice-Michaela IACOMI – Faculty of Agriculture, USAMV Bucharest, Romania
- Cristian IACOMI – Faculty of Agriculture, USAMV Bucharest, Romania
- Florin IMBREA – Faculty of Agriculture, USAMV of Banat Timișoara, Romania
- Viorel ION – Faculty of Agriculture, USAMV Bucharest, Romania
- Gheorghe JIGĂU – State University of Moldova, Republic of Moldova
- Ismail KARACA – Suleyman Demirel University, Turkey
- Mohammad KHANVAMI – Bu-Ali Sina University, Iran
- Gerard JITĂREANU – Faculty of Agriculture, USAMV Iași, Romania
- Doru-Ioan MARIN – Faculty of Agriculture, USAMV Bucharest, Romania
- Mircea MIHALACHE – Faculty of Agriculture, USAMV Bucharest, Romania
- Mihai NICOLESCU – Academy of Agricultural and Forestry Sciences, Romania
- Aurelian PENESCU – Faculty of Agriculture, USAMV Bucharest, Romania
- Françoise PICARD-BONNAUD – University of Angers, France
- Gheorghe Valentin ROMAN – Faculty of Agriculture, USAMV Bucharest, Romania
- Mihail RURAC – State Agrarian University of Moldova, Republic of Moldova
- Teodor RUSU – Faculty of Agriculture, USAMV Cluj-Napoca, Romania
- Dumitru Ilie SÂNDOIU – Faculty of Agriculture, USAMV Bucharest, Romania
- Philippe SIMONEAU – University of Angers, France
- Gheorghe SIN – Academy of Agricultural and Forestry Sciences, Romania
- Marin SOARE – Faculty of Agriculture and Horticulture, Craiova University, Romania
- Vasilica STAN – Faculty of Agriculture, USAMV Bucharest, Romania
- David C. WEINDORF – Louisiana State University AgCenter, USA

EDITORIAL BOARD

General Editor: Costică CIONTU

Executive Editor: Viorel ION

Members: Adrian Gheorghe BĂȘA, Lenuța Iuliana EPURE, André FALISSE, Leonard ILIE, Gheorghe JIGĂU, Doru Ioan MARIN, Mircea MIHALACHE

PUBLISHER:

**University of Agronomic Sciences and Veterinary Medicine of Bucharest,
Faculty of Agriculture, Romania**

Address: 59 Mărăști Blvd, District 1, 011464, Bucharest, Romania

Phone/Fax: + 40 213 318-0466; E-mail: journal@agro-bucuresti.ro

Webpage: <http://agronomyjournal.usamv.ro>

Copyright 2016

To be cited: Scientific Papers. Series A. Agronomy, Vol. LIX, 2016

The publisher is not responsible for the opinions published in the Volume. They represent the authors' point of view.

ISSN 2285-5785; ISSN CD-ROM 2285-5793; ISSN Online 2285-5807; ISSN-L 2285-5785

International Database Indexing: CABI, Index Copernicus, Google Scholar, CNCSIS B+, Ulrich's Periodicals Directory, Research Bible, Scipio, Scientific Indexing Service, PBN (Polish Scholarly Bibliography), OCLC (WorldCat)

SUMMARY

SOIL SCIENCES

1. **SOIL CO₂ EMISSIONS UNDER CONVENTIONAL AND CONSERVATIONAL TILLAGE METHODS IN SOYBEAN CULTIVATION IN CUKUROVA PLAIN OF TURKEY** - Davut AKBOLAT, Zeliha B. BARUT, M. Murat TURGUT, İsmail ÇELİK 15
2. **POTENTIAL OF THE LAND IN ARCHAR VILLAGE FOR CREATION OF VINES FOR HIGH-QUALITY WINE GRAPE VARIETIES. SOIL SPECIALITY OF THE TERROIR** - Mladen ALMALIEV, Krasimir TRENDAFILOV, Violeta VALCHEVA, Nedialka YORDANOVA, Nikolay MINEV 21
3. **PRODUCTIVITY ELEMENTS IN CONSERVATIVE AND CONVENTIONAL TILLAGE SYSTEMS** - Valentina ANDRIUCA, Ion BACEAN, Nicolai CAZMALI, Lucia MACRII, Rodica MELNIC 27
4. **QUALITY ASSESSMENT OF AN IRRIGATED FLUVISOL** - Valerian CERBARI, Gheorghe STEGARESCU 33
5. **AGROCHEMICAL STATE OF BROOMRAPE AFFECTED CHERNOZEMS OF CENTRAL AND SOUTH REGION OF THE REPUBLIC OF MOLDOVA** - Elena CERNOLEV, Adriana ACCIU, Olessea TABARA, Gheorghe JIGAU 39
6. **EFFECT OF SOIL WORKS ON CORN PRODUCTION IN SOUTH EAST BARAGAN** - Claudiu CHIȚOI, Costică CIONTU 45
7. **ANTROPIC DEGRADATION OF TYPICAL CHERNOZEMS AND THEIR RESTORATION UNDER STEPPE VEGETATION** - Tatiana CIOLACU 50
8. **EROSION - MAJOR PROBLEM IN THE EFFICIENT USE OF SLOPING LANDS** - Olessea COJOCARU, Esmeralda CHIORESCU 55
9. **STUDY OF THE BIOLOGICAL NITROGEN FIXATION PROCESS TO COMBINATION OF SOME BLUE-GREEN ALGAE FROM THE GENUS NOSTOC** - Sergiu DOBROJAN, Victor ȘALARU, Irina STRATULAT, Galina DOBROJAN 59
10. **CONTRIBUTION OF GREEN MANURE, RHIZOBIUM AND HUMIC + FULVIC ACID ON RECOVERING SOIL BIOLOGIC ACTIVITY OF OLIVE MILL WASTEWATER CONTAMINATED SOIL** - Kemal DOGAN, Ali SARIOGLU, Ali COSKAN 63
11. **EFFECT OF HUMIC + FULVIC ACID APPLICATION AT DIFFERENT DOSES ON BIOLOGICAL ACTIVITY OF DIFFERENT REGION SOILS** - Hande EROL, Ali COSKAN 69

12. RESEARCH ON THE IMPACT OF THE MORPHOLOGICAL VARIATION OF MINERAL FRAGMENTS ON SOIL HYDRAULIC PROPERTIES - Viorel FILER, Florian STĂTESCU, Madalin BODEA, Daniela RUSU, Florica DOROFTEI	75
13. THE PEDOFUNCTIONAL FRAMEWORK FOR ENSURING ADAPTIVE - LANDSCAPE - AMELIORATIVE TECHNOLOGIES - Gheorghe JIGAU, Elena TOFAN	83
14. THE BASIC FRAMEWORK FOR CONCEPTUAL-METHODICAL BIOENERGY RESOURCES OF CHERNOZEMS - Gheorghe JIGAU, Ana BARSAN, Elena TOFAN	89
15. THE INFLUENCE OF SOIL TILLAGE ON THE TEXTURE OF CHERNOZEM CAMBIC FROM MOLDOVA - Corina LEAH	96
16. GRASSLANDS OF MOLDOVA: QUALITY STATUS, VULNERABILITY TO ANTHROPOGENIC FACTORS AND ADAPTATION MEASURES - Tamara LEAH	100
17. COMPARATIVE EVALUATION OF PLASTICITY AND MICROAGGREGATES CONTENT OF CARBONATE CHERNOZEM UNDER DIFFERENT ANTHROPIC-IMPACT LEVELS - Lucia MACRII	106
18. RATIONAL METHODS OF SOILS MAINTENANCE IN ORCHARDS - Tatiana NAGACEVSCHI, Sava GRIȚCAN	110
19. DIVERSITY, DISTRIBUTION AND ECOLOGY OF THE FRESHWATER NATURAL HABITATS FROM SOUTHERN OF OLTENIA, ROMANIA - Mariana NICULESCU	116
20. NITRATES AND DRINKING WATER - Claudia Mariana ODOROG	122
21. EFFECT OF DIFFERENT PLASTIC COVER MATERIALS AND BIOFUMIGATION TO SOIL ORGANIC MATTER DECOMPOSITION IN GREENHOUSE SOLARIZATION - Hasan ÖZ, Ali COŞKAN, Atılğan ATILGAN	127
22. SOIL CHARACTERIZATION FROM OSOI - MORENI AREA, IAȘI, BY ANALYZING CERTAIN INDICATORS OF SALINITY - Maria Cătălina PASTIA, Florian STĂTESCU, Adriana Nicoleta STAN	130
23. BIOTA'S EVOLUTION OF GRAY FOREST SOILS IN THE CENTRAL ZONE OF THE REPUBLIC OF MOLDOVA - Irina SENICOVSCAIA	136
24. INFLUENCE OF SOIL TILLAGE UPON THE YIELD OF SEVERAL SEMI-LATE MAIZE HYBRIDS GROWN AT MOARA DOMNEASCĂ - ILFOV - Ionuț Cosmin SFETCU, Doru Ioan MARIN, Elena Loredana SFETCU	144
25. RESEARCH REGARDING THE USE OF GRAIN MASH AS SOIL FERTILIZER - Andrei SIURIS, Tatiana CIOLACU	149

26. A COMPARISON OF THE GRAVIMETRIC AND TDR METHODS IN TERMS OF DETERMINING THE SOIL WATER CONTENT OF THE CORN PLANT - Cagatay TANRIVERDI, Hasan DEGIRMENCI, Engin GONEN, Sedat BOYACI	153
27. POTENTIAL OF THE LAND IN ARCHAR VILLAGE FOR CREATION OF VINES FOR QUALITY WINE GRAPE VARIETIES. CLIMATIC AND GEOGRAPHIC SPECIALITY OF THE TERROIR - Krasimir TRENDAFILOV, Violeta VALCHEVA, Mladen ALMALIEV	159

CROP SCIENCES

1. SCREENING FOR ROOT AND SHOOT TRAITS OF DIFFERENT WHEAT GENOTYPES AND WILD RELATIVES UNDER FIELD CONDITIONS - Hayati AKMAN, Necdet AKGUN, Ahmet TAMKOC	167
2. CULTIVATION POTENTIAL OF <i>Salvia tomentosa</i> AND <i>S. aramiensis</i> UNDER THE EASTERN MEDITERRANEAN CONDITIONS - Mehmet ARSLAN	174
3. HERBAGE YIELD, ESSENTIAL OIL CONTENT AND COMPONENTS OF CULTIVATED AND NATURALLY GROWN <i>Origanum syriacum</i> - Mehmet ARSLAN	178
4. THE INFLUENCE OF GENOTYPE AND CLIMATIC FACTORS ON THE QUALITY OF SPRING BARLEY GROWN IN NORTH EAST BARAGAN - Nicoleta AXINTI, Maria CONTOMAN, Gabriela Alina CIOROMELE	183
5. POTATO TUBER SPROUT ROT CAUSED BY <i>Fusarium sambucinum</i> IN TURKEY - Mehmet Hadi AYDIN, Firat PALA, Cevdet KAPLAN	189
6. THE NPK FERTILIZATION EFFECTS OF TUBERS STARCH, DRY MATTER AND REDUCING SUGAR CONTENT - Nina BĂRĂSCU, Maria IANOȘI, Marcel M. DUDA, Edward MUNTEAN	194
7. STUDY OF NEMATODES ASSOCIATED WITH THE VEGETABLE CROPS IN SOME LOCALITIES IN TIARET (WEST OF ALGERIA) - Fatima Zohra BISSAAD, Farid BOUNACEUR	200
8. EFFECT OF <i>Bacillus thuringiensis</i> AND DIMILIN ON THE DATE MOTH <i>Ectomyelois ceratoniae</i> ZELLER - Fatima Zohra BISSAAD, Farid BOUNACEUR, Bahia DOUMANDJI-MITICHE	208
9. THE STUDY OF THE INFLUENCE OF THE LIQUID ORGANIC FERTILIZER ON THE PROCESS OF EMERGENCE AND DEVELOPMENT OF MAIZE - Tatiana BOCLACI, Larisa CREMENEAC	214

10. DISTRIBUTION, LIFE HISTORY AND BIOLOGY OF ALMOND SAWFLY (<i>Cimbex quadrimaculata</i> (Müller, 1766), Hymenoptera: Cimbicidae) - Halil BOLU	219
11. VARIOUS OBSERVATIONS ON SOME BIOLOGICAL CHARACTER OF PISTACHIO WHITE SCALE [<i>Suturaspis pistaciae</i> (Lindinger) Hemiptera: <i>Sternorrhyncha: Coccoidea: Diaspididae</i>] - Halil BOLU	223
12. THE YIELD AND GRAIN QUALITY OF WINTER TRITICALE VARIETY INGEN 93 IN THE MULTIFACTORIAL EXPERIMENT - Victor BURDUJAN, Mihail RURAC, Angela MELNIC, Victor STARODUB, Sergiu GAVRILAS, Vasile DREBOT	227
13. PRODUCTIVITY OF THE WINTER DURUM WHEAT IN POLYFACTO- RIAL EXPERIENCES - Victor BURDUJAN, Nicolae PALADE, Mihail RURAC, Victor STARODUB, Angela MELNIC, Sergiu GAVRILAŞ	231
14. THE USE OF ENTOMOPATHOGENS IN THE CONTROLLING OF INSECT PESTS OF STORED PRODUCT - Ramazan CANHILAL	235
15. ECONOMIC THRESHOLD IN THE MANAGEMENT OF THE SUNN PEST - Ramazan CANHILAL	241
16. STEINERNEMATID AND HETERORHABDITID NEMATODES AGAINST <i>Zabrus</i> SPP. - Ramazan CANHİLAL	247
17. THE EFFECTS OF DIFFERENT HARVEST TIMES ON FORAGE YIELD AND QUALITY OF SOME VETCH (<i>Vicia</i> spp.) SPECIES - Ihsan ÇETIN, Mevlut TURK	251
18. THE NEW WAY OF EVALUATING THE PROTEIN POLYMORPHISM FOR MAIZE BREEDING AND SEED PRODUCTION - Galina COMAROVA, Alexandr ROTARI, Eugen ROTARI	257
19. STABILITY VALUATION OF SOME MIXTURES BETWEEN RETARDANTS AND ANTIBROADLEAVED HERBICIDES FOR THE GRAIN YIELD OF DURUM WHEAT - Grozi DELCHEV	261
20. STABILITY VALUATION OF SOME MIXTURES BETWEEN FOLIAR FERTILIZERS AND ANTIGRAMINACEOUS HERBICIDES FOR THE GRAIN YIELD OF DURUM WHEAT - Grozi DELCHEV	267
21. THE EFFICIENCY USE OF DOUBLED-HAPLOID TECHNOLOGY IN MAIZE BREEDING – OBTAINING DH PARENT LINES AND HYBRIDS - Georgeta DICU, Stelica CRISTEA	273
22. INFLUENCE OF FERTILIZATION ON RYE PRODUCTION IN THE SANDY SOILS CONDITIONS FROM SOUTHERN OLTENIA - Iulian DRĂGHICI, Reta DRĂGHICI, Mihaela CROITORU	279
23. THE STUDY OF SOME CULTIVARS OF COWPEA UNDER CLIMATE CHANGE IN SOUTHERN OLTENIA - Reta DRĂGHICI, Iulian DRĂGHICI, Mihaela CROITORU	283

24. INFLUENCE OF DIFFERENT LEVELS OF WATER SUPPLY ON PRODUCTION AND ECONOMIC EFFICIENCY UNDER MAIZE CULTIVATED IN DOBROGEA REGION OF ROMANIA - Cristian Laurentiu DRAGOMIR, Elena PARTAL	289
25. ACHIEVEMENTS AND PROBLEMS IN THE WEED CONTROL IN BARLEY (<i>Hordeum vulgare</i> L.) - Mitko GEORGIEV, Grozi DELCHEV	294
26. MEDICINAL PLANT CROPS - IMPORTANT SOURCE OF HIGH VALUE-ADDED PRODUCTS - Alice GRIGORE, Lucia PIRVU, Corina BUBUEANU, Svetlana COLCERU-MIHUL, Carmen IONITA, Lucian IONITA	298
27. ASSESSMENT OF AQUACROP MODEL IN THE SIMULATION OF WHEAT GROWTH UNDER DIFFERENT WATER REGIMES - Sema KALE	308
28. IDENTIFICATION OF PROSTIGMATID MITES ASSOCIATED WITH HAWTHORN TREES IN WEST OF IRAN - Masoumeh KHANJANI, Mohammad KHANJANI, Fatemeh AMINI	315
29. EFFECT OF NUTRIENTS SUPPLY WITH FOLIAR APPLICATION ON GROWING DEGREE DAYS, PROTEIN AND FATTY YIELD OF CORN IN MEDITERRANEAN CONDITIONS - Yakup Onur KOCA	318
30. EFFECT OF NITROGEN ON THE TRANSLOCATION OF DRY MASS AND NITROGEN IN BARLEY - Svetla KOSTADINOVA, Galia PANAYOTOVA, Lyubena KUZMANOVA	327
31. IRRIGATION SCHEDULING AND THE IMPACT OF IRRIGATION ON THE YIELD AND YIELD COMPONENTS OF SWEET CORN - Milena MOTEVA, Nelly GADJALSKA, Viktoria KANCHEVA, Todor TASHEV, Veska GEORGIEVA, Natalia KOLEVA, Ivan MORTEV, Vesela PETROVA-BRAHICHEVA	332
32. INFLUENCE OF BASIC TILLAGE DEPTH ON SOIL HUMIDITY RESOURCES AND SUNFLOWER YIELDS - Angelina MUHOVA, Stefka DOBREVA, Ivan SALDZHIEV	340
33. STUDY REGARDING ORGANIC AGRICULTURE AND CERTIFICATION OF PRODUCTS - Paula Ionela NASTASE, Maria TOADER	344
34. IDENTIFICATION AND CHARACTERIZATION OF MUTANTS INDUCED BY GAMMA RADIATION IN WINTER WHEAT (<i>Triticum aestivum</i> L.) - Mykola NAZARENKO	350
35. THE RESPONSE TO STOMATAL CONDUCTANCE AND CHLOROPHYLL VALUE OF GROUNDNUT GROWN UNDER SALT STRESS IN DIFFERENT DEVELOPMENT STAGES - Berkant ÖDEMİŞ	354
36. ANTIFUNGAL ACTIVITY OF <i>Liquidambar orientalis</i> L., and <i>Myrtus communis</i> L. AGAINST SOME PLANT PATHOGENIC FUNGI - Abdurrahman ONARAN, Yusuf BAYAN	360

37. DOWNY MILDEWS SPECIES ON THE WEEDS OF LENTIL FIELDS IN DIYARBAKIR IN TURKEY - Cumali ÖZASLAN	365
38. INFLUENCE OF NITROGEN FERTILIZATION ON BULGARIAN COTTON CULTIVARS - Galia PANAYOTOVA, Svetla KOSTADINOVA, Neli VALKOVA	368
39. HERBAGE YIELD AND QUALITY OF WHEAT STUBBLE AND SORGHUM SUDAN-GRASS PASTURES - Altngül Özasan PARLAK, Ahmet GÖKKUŞ, Firat ALATÜRK, Hülya HANOĞLU, Cemil TÖLÜ	374
40. APPLYING OF MANURE TO WINTER WHEAT ÎN BESSARABIA - Gheorghe PRIPA	378
41. OLD AND REGIONAL VARIETIES OF PEAR TREES IN THE TERRITORY OF CZECH SILESIA - Radim LOKOČ, Maria TOADER	382
42. CONTRIBUTIONS TO THE ESTABLISHMENT OF THE FERTILIZATION SYSTEM WITH MINERAL NITROGEN FOR AUTUMN CROPS RAPE AND WHEAT - IN THE SPECIFIC ENVIRONMENT OF BULBUCATA GIURGIU - Luxița RÎȘNOVEANU, Liviu DINCĂ, Emil Igor Vlad GEORGESCU, Mariana Carmen BURTEA	390
43. EFFECTS OF DIFFERENT PHOSPHORUS LEVELS ON UREIDES CONTENTS AND NODULATION OF SOYBEAN (<i>Glycine max</i> L.) IN RELATION TO SOIL MOISTURE REGIME - Vladimir ROTARU	397
44. DETERMINATION OF OPTIMUM ROW-SPACING AND PLANT DENSITY IN GOLDASHTI SAFFLOWER VARIETY - Mohammad SHARIF MOGHADDASI, Amir Hasan OMIDI	403
45. THE STUDY OF THE DROUGHT TOLERANCE OF CANOLA (<i>Brassica napus</i> L.) CULTIVARS BY USING STRESS TOLERANCE INDICES - Seyed Habib SHOJAEI, Khodadad MOSTAFAVI, Pooria MAZLOOM, Peyman KERMANI, Sajjad SHOJAEI, Seyede Mitra SADATI	409
46. EFFECT OF SOWING TIME ON PEANUT (<i>Arachis hypogaea</i> L.) CULTIVARS: I. YIELD, YIELD COMPONENTS, OIL AND PROTEIN CONTENT - Tahsin SOGUT, Ferhat OZTURK, Suleyman KIZIL	415
47. NUTRITION VALUE OF TWO GRAIN COMMON WHEAT - Antoniya STOYANOVA, Gancho GANCHEV, Velika KUNEVA	421
48. IMPACT OF SETTLEMENT POINTS ON SOME PROPERTIES OF A HIGHLAND RANGELAND VEGETATION OF EASTERN ANATOLIA REGION OF TURKEY - Mustafa SÜRMEŒ, Ali KOÇ	426
49. PROJECTIONS OF CHANGES IN PRODUCTIVITY OF MAJOR AGRICULTURAL CROPS IN THE REPUBLIC OF MOLDOVA ACCORDING TO CMIP5 ENSEMBLE OF 21 GCMs FOR RCP2.6, RCP4.5 AND RCP8.5 SCENARIOS - Lilia ȚĂRANU	431

50. AGROBIOLOGICAL PECULIARITIES AND PROSPECTS OF THE <i>Lathyrus perennial</i> SPECIES IN MOLDOVA - Alexandru TELEUȚĂ, Victor ȚÎȚEI	441
51. SPECIES OF PERENNIAL GRASSES AS FEEDSTOCK FOR BIOGAS PRODUCTION IN THE REPUBLIC OF MOLDOVA - Victor ȚÎȚEI	447
52. ESTABLISHING SOWING DENSITY AND OPTIMAL DOSAGE OF FERTILIZER FOR INCREASING PRODUCTION OF WINTER WHEAT AND BARLEY, UNDER CLIMATE CHANGE CONDITIONS - Daniela TRIFAN, Marcel BULARDA, Radu ISPAS, Nicolae POPESCU	453
53. EFFECTS OF FERTILIZATION ON SEED YIELD AND FORAGE QUALITY OF COMMON VETCH (<i>Vicia sativa</i> Roth.) - Mevlüt TÜRK, Fatma YILDIZ	462
54. EFFECTS OF SALT STRESS ON GERMINATION OF SOME SILAGE MAIZE (<i>Zea mays</i> L.) CULTIVARS - Mevlüt TURK, Ozge ESER	466
55. MINIMIZING OF SIDE EFFECTS OF PESTICIDES ON AGRICULTURAL ENVIRONMENT - Nigar YARPUZ-BOZDOGAN, Ali Musa BOZDOGAN	470
56. VEGETATION FEATURES OF ALPINE AND SUBALPINE RANGELANDS IN EASTERN BLACK SEA REGION - Tamer YAVUZ, Mustafa SÜRMEŃ	474
57. PRODUCTIVITY AND YIELD STABILITY OF SIX GRAIN LEGUMES IN THE MODERATE CLIMATIC CONDITIONS OF BULGARIA - Tsenka ZHELIAZKOVA, Dimitar PAVLOV, Grosi DELCHEV, Antoniya STOYANOVA	478

MISCELLANEOUS

1. MIXING AMELIORANTS IN SOIL WITH DIFFERENT ACTIVE DISK WORK BODIES - Manol DALLEV, Ivan IVANOV, Ivan MITKOV, Jordanka ZAPRJANOVA	491
2. SYSTEM FOR RECORDING THE ANALYSIS RESULTS AS PART OF THE QUALITY CONTROL PLANES IN A STARCH FACTORY - Georgiana DECIU, Arina Oana ANTOCE	496
3. INSECTICIDES IMPACT ON SEED GERMINATION AND EARLY SEEDLINGS GROWTH IN MAIZE (<i>Zea mays</i> L.) - Elena DELIAN, Elena SĂVULESCU, Ion SĂVULESCU, Adrian Marius TEBAN	503
4. INFLUENCE OF DRYING PROCESS ON PHENOLIC CONTENT AND ANTIOXIDANT ACTIVITY OF TWO DIFFERENT AUTOCHTHONOUS ALBANIAN FIG VARIETIES - Luziana HOXHA, Renata KONGOLI	509
5. THE PARADOX OF MALATHION USED FOR HUMAN LIFE - Yakup Onur KOCA, Ahmet Onder USTUNDAG, Melis YALCIN	515

6. AN INVESTIGATION INTO THE RAM EFFECT ON SYNCHRONISING OESTRUS AND IMPROVING CONCEPTION AND PREGNANCY RATES OF A HILL FLOCK DURING THE BREEDING SEASON - Carol MOFFAT, Anya WESTLAND, Lancelot BUTTERS	521
7. DETERMINATION OF BIOGAS POTENTIAL OF AYDIN PROVINCE - Berker ÖZTÜRK, Ahmet KILIÇKAN	531
8. MORPHO-ANATOMY AND ADAPTATION TO SOME ROMANIAN AQUATIC ENVIRONMENTS OF <i>Nymphoides peltata</i> (GMEL.) O. KUNTZE (<i>Asterales: Menyanthaceae</i>) - Emilia Brîndușa SĂNDULESCU, Gina VASILE SCĂEȚEANU, Tudor ȘCHIOPU, Nicoleta OLTENACU, Mala-Maria STAVRESCU-BEDIVAN	537
9. PROSPECTS FOR THE UTILIZATION OF THE <i>Miscanthus giganteus</i> AND <i>Polygonum sachalinense</i> FOR SOLID BIOFUEL PRODUCTION IN THE REPUBLIC OF MOLDOVA - Victor ȚÎȚEI, Andrei GUDÎMA, Alexandru MUNTEAN, Mihai GADIBADI, Aurelia LUPAN	543
10. DETERMINATION OF PESTICIDE RESIDUES IN TOMATOES COLLECTED FROM AYDIN PROVINCE OF TURKEY - Melis YALÇIN, Cafer TURGUT	547
11. DETERMINING THE LOCATION OF THE ASSEMBLY CENTER FOR EXTRUSION THE WASTE FROM ROSE PRODUCTION IN REPUBLIC OF BULGARIA - Ivan ZAHARIEV, Dimitar KEHAYOV	552