INFLUENCE OF DIFFERENT *IN VITRO* SIMULATORS FOR HYDRIC STRESS FOR GROWTH AND DEVELOPMENT OF POTATO

Andreea TICAN¹, Mihaela CIOLOCA¹, Nicoleta CHIRU¹, Carmen BĂDĂRĂU^{1,2}

¹National Institute of Research and Development for Potato and Sugar Beet Brasov, 2 Fundaturii Street, Brasov, Romania
²Faculty of Food and Tourism, Transilvania University of Brasov, 148 Castelului Street, Brasov Romania

Corresponding author email: tican_andreea@yahoo.com

Abstract

Necessity of finding genotypes adapted to drought has become urgent due to the effect of this type of stress on potato production. The most important phase, indispensable for improving drought tolerance is to identify genotypes tolerant and sensitive to drought. In this study, to induce in vitro water stress were used polyethylene glycol and sorbitol that was comparable with the basic medium MS, considered control. Determinations were performed 4 weeks after inoculation of mini cuttings belonging to five varieties of plantlets (Ruxandra, Sarmis, Gared, Marvis, Rustic) and the parameters analyzed were next: number of leaves, number of internodes, height of plantlets, root length, weight of fresh plantlet, weight of fresh root. Medium in which was added PEG with different concentrations significantly reduced the average weight of fresh plantlet and root compared with the control medium and medium with sorbitol and significantly reduced the mean number of internodes, the average height of the plantlet, the average root length. This osmotic agent (PEG) can be recommended for in vitro simulation of drought to identify tolerant genotypes to hydric stress.

Key words: hydric stress, in vitro, osmotic agent, potato.

INTRODUCTION

Water deficit, extreme temperatures and low atmospheric humidity lead to drought, which is one of the limiting factors affecting crop quality. The amplitude of the effects of drought on potato production depends on phenological calendar, duration and severity of stress. Two critical periods: sprouting and tuberization affect in final tubers production. Potato crop is often considered sensitive to drought and production during successive episodes of drought may be compromised. Necessity to identify genotypes adapted to drought has become urgent because of the effect this type of stress on the growth of potato and production of this crop. Particularly important for production of potato tubers is the precipitation amount during the growing season and the distribution on the vegetative stages. It is estimated that during the growing season are required 250-400 mm precipitations (Bîlteanu, 2001).

The ability of roots to penetrate the soil depends on the power that can exert roots and may be associated with drought tolerance (Tardieu, 1994). If plants cannot take up water

from the soil needed to compensate for the lost through perspiration, install wilting phenomenon, a consequence of the drought effects (Ianoşi, 2002).

Drought reduces the growth of roots, at the time of sprouting. Drought installed after plant sprouting inhibits the stolons development, thereby reducing the number of tubers. These processes are irreversible, even though soil moisture subsequently recovers. Capacity of roots to penetrate into the soil depends on the power that roots can exert and may be associated with drought tolerance.

In vitro tissue culture allowed a deeper understanding of the physiology and biochemistry of plants grown under unfavorable environmental (Benderradji L. and colab., 2012).

The most widely used method for the selection of genotypes tolerant to abiotic stress is the *in vitro* selection pressure technique. This is based on the *in vitro* culture of plant cells, tissues or organs on a medium supplemented with selective agents, allowing selecting and regenerating plants with desirable characteristics (Pérez-Clemente and Gómez-Cadenas, 2012). Polyethylene glycol (PEG), sucrose, mannitol or sorbitol have been used by several workers as osmotic stress agents for *in vitro* selection.

In vitro simulation drought was made to identify varieties with optimum tolerance at drought. Observations showed that was obtained a slowing of regeneration cuttings. Sensitivity to drought was not uniform for varieties analyzed. For in vitro selection PEG. sucrose, mannitol and sorbitol were analyzed in several research papers, as agents of osmotic stress. Sorbitol is a sugar alcohol hexahvdrate with osmotic effect. Widely water stress in vitro simulation is used polyethylene glycol. Culture media which contain PEG, imitate dry soil, rather than the culture media which have low molecular weight compounds. With the increasing amount of sorbitol and PEG, water absorption becomes difficult for plantlets from nutrient medium and thus is simulate the effect of drought. *In vitro* culture technique minimizes external environmental variations due to nutrient medium defined and controlled conditions and homogeneity of stress applied.

MATERIALS AND METHODS

In Laborator of Vegetal Tissue Culture, of NIRDPSB Brasov (2015) was made a study for identy the adequate agente for induce water stress. Microplantlets from the culture collection were multiplied to each internode cuttings and were inoculated and on Murashige-Skoog medium. Plantlets which were developed were multiplied to obtain nodal cuttings. Cuttings of these plantlets were used as explants, for further multiplication in vitro. As a basic medium was used Murashige-Skoog medium, naphthylacetic acid, sucrose, agar (this was considered control medium). Both PEG and sorbitol were added in the culture medium in 4 concentrations (0.5%; 1.0%; 1.5%; 2.0%). Test tubes with mini cuttings of varieties proposed for in vitro water stress test, were placed in the growth chamber by ensuring light and temperature regime required for growth and development of plantlets. After 4 weeks determinations were made for analysis of the following parameters, for a part of plantlets of these varieties: number of leaves and internodes / plantlet and plantlet height, root length, fresh weight of plantlet and root

(figures 1, 2, 3, 4, 5). The results were processed by analysis of variance and the significance of differences was determined using the method of multiple comparisons, respectively Duncan test. Experimental differences higher than 5% are considered significant (Săulescu and Săulescu, 1967).

For determine the effect of water stress on developing plantlets the study consisted in an bifactorial experience of two factors (5 x 3), 4 repetitions, including the following factors:

Experimental factor A: variety, with 5 graduations:

- a1- Ruxandra;
- a₂- Sarmis;
- a₃- Gared;
- a₄- Marvis;
- a₅- Rustic.

Experimental factor B – nutrient media used with 3 graduations:

- b₁-control medium MS, to which was made no addition of osmotic agent;

- b₂- MS medium, to which was added PEG;
- b₃- MS medium, to which was added sorbitol.

RESULTS AND DISCUSSIONS

Treatments performed with PEG significantly reduced the average weight of plantlet and of root and compared to sorbitol and nutrient medium, considered control (MS) and significantly reduced the average number of internodes, the height of plantlet, root length. From Table 1 it is noted that PEG and sorbitol have no negative influence on the average number of leaves, but PEG reduces the average number of internodes compared to control medium.

For the first element in the study, Duncan test analysis indicates a proximity of values, with no significant differences between the number of leaves formed on MS medium (8.8 leaves) and on medium that contained sorbitol 1% (8.4 leaves), so compared with PEG, sorbitol has a less stressful effect on plantlets, there is a stronger competition in formation of plant leaves depending on the concentration (Table 2). PEG on maximum concentration of 2% had as result hydric stres for leaves formatic, these are in lowest number (7.40).



Fig. 1. Effects of water stress simulators (PEG and sorbitol) on growth and development of plantlets and roots for Ruxandra variety



Fig. 2. Effects of water stress simulators (PEG and sorbitol) on growth and development of plantlets and roots for Sarmis variety



Fig. 3. Effects of water stress simulators (PEG and sorbitol) on growth and development of plantlets and roots for Gared variety



Fig. 4. Effects of water stress simulators (PEG and sorbitol) on growth and development of plantlets and roots for Marvis variety



Fig. 5. Effects of water stress simulators (PEG and sorbitol) on growth and development of plantlets and roots for Rustic variety

The average number of internodes / plant indicates the effect of hydric stress attenuation for PEG, on minimum concentration of 0.5% (5.30 internods). Analysis of Duncan test do not record significant differences compared to this value for nutrient medium on which sorbitol was used at concentrations 0.5; 1; 1.5% (BC). Sorbitol 2% is so drastic as 1 and 1.5% PEG (Duncan test D and CD) for inducing *in vitro* drought.

The average height of plantlets (cm) is similarly influenced by the first concentration of two chemicals of water stress accelerators (7.725 and 7.685 respectively) (B), while the last two concentrations studied, indicating a similar effect, but severely on growth *in vitro* plantlets (5.515 cm, with decreasing at 3.645 cm for PEG and respectively 5.29 cm, with decreasing to 3.685 cm by application of sorbitol).

A very decisive indicator of drought simulation in vitro is root length (cm). Concentrations of 1 and 1.5% for sorbitol substance does not describe a relevant effect of water stress (there are no significant differences) with values close the control medium (7.030 cm).

The concentration of 0.5% sorbitol is less conclusive, because should be expected to induce less attenuating stress for increasing root length of microplant, but this concentration results in the formation of a roots with an average length less (6.15 cm), but with a lower average weight close to that obtained using the control medium.

For concentration of 0.5% sorbitol, it would be expected to result in a less pronounced stress on the plant, on the increase in root length, but in fact lead shortening roots, but with a value of average root weight (79,275mg) (B), close to the value obtained by using of control medium (89,595mg) (A).

PEG-induced very well drought at concentrations of 1.5 and 2%; produces very little roots with weight of 18.610 mg 23.785 mg (Duncan test G, H).

From Table 3 Marvis variety combat drought effect by forming a large number of leaves (with an average of 9.083 leaves). This is followed by Rustic variety (8.222 leaf / plantlet). The average number of internodes grouped in a closely way Marvis, Ruxandra and Rustic varieties (A), followed by Gared and Sarmis varieties (B). For plant height significantly detaches Ruxandra variety which registered a value of 7.269 cm (A), followed by Sarmis variety, which registered 6.892 cm (B). The root has a higher resistance to drought than the plant. It is noted in this sense Ruxandra and Gared varieties, with high and close values (7.111 cm; 7.006 cm) (A), followed by Marvis variety (5.758 cm). On the opposite side the

Sarmis variety had a tendency of slow root growth (only 4.6222 cm). For fresh plantlet weight (mg) is distinguished Ruxandra variety (126.142 mg), followed by Rustic variety (120.397 mg).

Gared variety, which in terms of microplants size is located on last place (with an average height of 5.506 cm), also recorded an average low weight of microplant 92.019 mg (situated on the last place). Regarding the average weight of fresh root (mg) detaches Ruxandra variety, with a very high value of 83.9 mg (A). followed by Gared (50,489 mg) and Sarmis variety (48,986). Regarding to this variety from previous analyzes, it appears that shows a tolerance to hydric stress, by producing microplants with a high height (6.892 cm) and with a high number of internodes (4.556). Rustic variety take last place, form roots with an average length small (5.244 cm) and a low weight (34.256 mg) indicating an inability of variety to fight with in vitro hydric stress.

Table 1 - Simulators osmotic stress influence compared to the control medium for the elements of growth and development for microplants

| The treatment made | Average number of leaves | Average number of internodes | The average height of plantlets (cm) | The average length of root (cm) | The average weight of fresh plantlet (mg) | The average root weight (mg) |
|------------------------|---|------------------------------------|---|---------------------------------|---|---------------------------------|
| Control medium (MS) | 8.800 A | 6.200 A | 10.70 A | 7.025 A | 207.535 A | 89.595 A |
| MS+PEG | 7.713 A | 4.663 B | 5.708 B | 5.166 B | 86.626 C | 35.750 C |
| MS+sorbitol | 8.013 A | 4.950 AB | 5.779 B | 6.711 AB | 99.928 B | 60.750 B |
| Med | Media that we are in the same column followed by the same letters are not significant, according to Duncan test | | | | | |

| LSD=1.246 | LSD=1.416 | LSD=1.812 cm | LSD=1.252 cm | LSD=9.386 mg | LSD=4.865 mg |
|-----------|-----------|--------------|--------------|--------------|--------------|

Table 2 - Influence of osmotic stress concentrations simulators on the elements of growth and development of microplants

| Hydric stress simulator concentration (%) | Average number of leaves | Average number of internodes | The average height of plantlets (cm) | The average length of root (cm) | The average weight of fresh plantlet (mg) | The average root weight (mg) |
|---|---|--|--|--|--|--|
| - | 8.80 A | 6.20 A | 10.70 A | 7.030 A | 207.535 A | 89.595 A |
| 0.5 | 7.95 CD | 5.30 B | 7.725 B | 6.085 C | 144.200 C | 56.235 C |
| 1 | 7.65 DE | 4.55 D | 5.945 D | 5.025 D | 83.285 F | 44.370 E |
| 1.5 | 7.85 CD | 4.90 CD | 5.515 E | 4.825 D | 71.245 G | 23.785 G |
| 2 | 7.40 E | 3.90 E | 3.645 F | 4.730 D | 47.775 H | 18.610 H |
| 0.5 | 7.75 CDE | 5.15 BC | 7.685 B | 6.150 C | 153.085 B | 79.275 B |
| 1 | 8.10 BC | 5.05 BC | 6.455 C | 6.835 A | 118.695 D | 78.275 B |
| 1.5 | 8.40 AB | 5.05 BC | 5.29 E | 7.210 A | 88.445 E | 54.360 D |
| 2 | 7.80 CDE | 4.55 D | 3.685 F | 6.650 B | 39.485 I | 31.090 F |
| _ | simulator concentration (%) - 0.5 1 1.5 2 0.5 1 1.5 2 2 0.5 1 1.5 2 2 | Initiation Initiation simulator number of leaves - 8.80 A 0.5 7.95 CD 1 7.65 DE 1.5 7.85 CD 2 7.40 E 0.5 7.75 CDE 1 8.10 BC 1.5 8.40 AB 2 7.80 CDE | Interaction (%) Interaction (%) simulator concentration (%) number of leaves . 8.80 A 6.20 A 0.5 7.95 CD 5.30 B 1 7.65 DE 4.55 D 1.5 7.85 CD 3.90 E 0.5 7.75 CDE 5.15 BC 1 8.10 BC 5.05 BC 1.5 8.40 AB 5.05 BC 2 7.80 CDE 4.55 D | Initializion concentration (%) Initializion number of leaves Initializion number of internodes Initializion plantlets (cm) - 8.80 A 6.20 A 10.70 A 0.5 7.95 CD 5.30 B 7.725 B 1 7.65 DE 4.55 D 5.945 D 1.5 7.85 CD 4.90 CD 5.515 E 2 7.40 E 3.90 E 3.645 F 0.5 7.75 CDE 5.15 BC 7.685 B 1 8.10 BC 5.05 BC 6.455 C 1.5 8.40 AB 5.05 BC 5.29 E 2 7.80 CDE 4.55 D 3.685 F | Internet Internet Internet simulator number of number of height of length of root - 8.80 A 6.20 A 10.70 A 7.030 A 0.5 7.95 CD 5.30 B 7.725 B 6.085 C 1 7.65 DE 4.55 D 5.945 D 5.025 D 1.5 7.85 CD 3.90 E 3.645 F 4.730 D 0.5 7.75 CDE 5.15 BC 7.685 B 6.150 C 1 8.10 BC 5.05 BC 6.455 C 6.835 A 1.5 8.40 AB 5.05 BC 5.29 E 7.210 A 2 7.80 CDE 4.55 D 3.685 F 6.650 B | Interlige Interlige Interlige Interlige Interlige Interlige Interlige simulator number of number of height of respective length of root weight of fresh - 8.80 A 6.20 A 10.70 A 7.030 A 207.535 A 0.5 7.95 CD 5.30 B 7.725 B 6.085 C 144.200 C 1 7.65 DE 4.55 D 5.945 D 5.025 D 83.285 F 1.5 7.85 CD 4.90 CD 5.515 E 4.825 D 71.245 G 2 7.40 E 3.90 E 3.645 F 4.730 D 47.775 H 0.5 7.75 CDE 5.15 BC 7.685 B 6.150 C 153.085 B 1 8.10 BC 5.05 BC 5.29 E 7.210 A 88.445 E 2 7.80 CDE 4.55 D 3.685 F 6.650 B 39.485 I |

Media that we are in the same column followed by the same letters are not significant, according to Duncan test

LSD=0.4218 LSD=0.3740 LSD=0.3778 cm LSD=0.4015 cm

LSD=12.277mg LSD=1.783mg

| Variety | Average number of leaves | Average number of internodes | The average height of plantlets (cm) | The average length of root (cm) | The average weight of fresh plantlet (mg) | The average root weight (mg) |
|----------|--------------------------------|------------------------------------|--|---------------------------------------|---|------------------------------------|
| Ruxandra | 7.528 CD | 5.194 A | 7.269 A | 7.111 A | 126.142 A | 83.9 A |
| Sarmis | 7.222 D | 4.556 B | 6.892 B | 4.6222 D | 95. 925 C | 48.986 C |
| Gared | 7.778 C | 4.667 B | 5.506 D | 7.006 A | 92.019 D | 50.489 B |
| Marvis | 9.083 A | 5.333 A | 5.700 D | 5.758 B | 95.378 C | 46.572 D |
| Rustic | 8.222 B | 5.056 A | 6.103 C | 5.244 C | 120.397 B | 34.256 E |
| Media t | that we are in the sa | me column followe | ed by the same letters a | are not significant, acc | ording to Duncan test | |

Table 3 - Behaviour the varieties tested under the influence of treatments made to induce water stress

LSD=0.3266 LSD=0.3183 LSD=0.3229 cm LSD=0.3638 cm LSD=1.9722 mg LSD=1.458 mg

CONCLUSIONS

Medium with different concentrations in which was additional PEG significantly reduced the weight of fresh plantlets and fresh root as compared with the control medium and sorbitol and also significantly reduced the number of internodes, the height of the plantlet, the root length. The smallest influence of water stress on plantlet height was observed for Ruxandra variety. By using PEG in culture medium, potato plantlets reach 7.83 cm. The roots are the primary sensors of water deficit. Gared and Ruxandra varieties showed a good tolerance to water stress for root length, by applying sorbitol on nutritive medium, plantlet root reaching an average length of 8.13 and 7.22 cm. By addition of PEG, it appears that the same varieties Ruxandra and Gared, shows tolerance to water stress, but PEG being an agent stronger the levels were lower.

REFERENCES

- Benderradji L., Brini F., Kellou K., Ykhelf N., Djekoun A., Masmoudi K., Bouzerour. H., 2012. Callus induction, proliferation, and plantlets regeneration of two bread wheat (*Triticum Aestivum* L.) genotypes under saline and heat stress conditions. ISRN Agronomy, Article ID 367851. Bobr. Plant Cell Tissue and Organ Culture; 102 387–395.
- Bîlteanu Gheorghe, 2001. Fitotehnie, Volumul 2, Editura Ceres, București.
- Ianoși, S. I., 2002. Bazele cultivării cartofului pentru consum, Editura Phoenix, Brașov.
- Pérez-Clemente R. M and Gómez-Cadenas A., 2012. In vitro Tissue Culture, a Tool for the Study and Breeding of Plants Subjected to Abiotic Stress Conditions. In: Annarita Leva and Laura M. R. Rinaldi(Eds.) Agricultural and Biological Sciences. "Recent Advances in Plant in vitro Culture", Chapter 5, 91-108.
- Săulescu, N.A, Săulescu, N.N., 1967. Cîmpul de experiență, Întreprinderea poligrafică Sibiu.
- Tardieu F., 1994. Growth and functioning of roots and of root systems subject to compaction towards a system with multiple signalling. Soil and Tillage Research 30, 217–43.

FOOD BIOTECHNOLOGY

SCIENTIFIC BULLETIN SERIES F. BIOTECHNOLOGIES Volume XX, 2016

University of Agronomic Sciences and Veterinary Medicine of Bucharest Faculty of Biotechnologies

SCIENTIFIC BULLETIN SERIES F. BIOTECHNOLOGIES VOLUME XX

2016 BucharesT

SCIENTIFIC COMMITTEE

- Veronika ABRAM Biotechnical Faculty, University of Ljubljana, Slovenia
- Petru ALEXE Faculty of Food Science and Engineering, University of "Dunarea de Jos Galati", Romania
- Ioan ARDELEAN Institute of Biology, Romanian Academy
- Narcisa BABEANU Faculty of Biotechnologies, UASVM Bucharest, Romania
- Gabriela BAHRIM Faculty of Food Science and Engineering, University of "Dunarea de Jos Galati", Romania
- Gustavo V. BARBOSA-CANOVAS Washington State University Pullman, State of Washington, US
- Judith BARETTE Manchester Metropolitan University, United Kingdom
- · Ingrid BAUMAN Faculty of Food Technology and Biotechnology, Zagreb, Croatia
- Nastasia BELC Faculty of Biotechnologies, UASVM Bucharest, Romania
- Daniela BORDA Faculty of Food Science and Engineering, University of "Dunarea de Jos Galati", Romania
- Dorica BOTAU Faculty of Agriculture, UASVM Banat, Timisoara, Romania
- Calina Petruta CORNEA Faculty of Biotechnologies, UASVM Bucharest, Romania
- Delia DIMITRIU Manchester Metropolitan University, United Kingdom
- Paulo Jose do AMARAL SOBRAL Depto de Eng. De Alimentos FZEA USP, Pirassununga, Brazil
- Katherine FLYNN European Association for Food Safety, Brussels, Belgium
- Helmut GLATTES ISEKI Food Association, Austria
- Gustavo Fidel GUTIERREZ-LOPEZ ENCB-IPN, National School of Biological Sciences, National Polytechnic Institute, Mexico
- Florentina ISRAEL ROMING- Faculty of Biotechnologies, UASVM Bucharest, Romania
- Stefana JURCOANE Faculty of Biotechnologies, UASVM Bucharest, Romania
- Huub LELIEVELD GHI Association Netherlands and EFFoST Executive Committee, Netherlands
- Florentina MATEI Faculty of Biotechnologies, UASVM Bucharest, Romania
- Lynn McINTYRE Food Microbiology, Harper Adams University College Edgmond, Newport, United Kingdom
- Amalia Carmen MITELUT Faculty of Biotechnologies, UASVM Bucharest, Romania
- Dumitru MILITARU Institute Pasteur, Bucharest, Romania
- Anca NICOLAU Faculty of Food Science and Engineering, University of "Dunarea de Jos Galati", Romania
- Tatiana VASSU University of Bucharest, Romania
- Estela de Oliveira NUNES Santa Catarina West University UNOESC Biotechnological Nucleus, Brazil
- Paola PITTIA Dipartimento di Scienze, degli Alimenti University degli Studi di Teramo, Italy
- Mona Elena POPA Faculty of Biotechnologies, UASVM Bucharest, Romania
- Cristina SILVA ISEKI Food, Catholic University of Portugal
- Margarida VIEIRA Directora do Dep. De Engenharia Alimentar, Instituto Superior de Engenharia, Universidade do Algarve, Portugal
- Medana ZAMFIR Institute of Biology, Romanian Academy

EDITORIAL BOARD

General Editor: Călina Petruța CORNEA

Executive Editor: Ștefana JURCOANE

Secretariat: Silvana DĂNĂILĂ-GUIDEA, Mira TURTOI, Luminița VIȘAN

PUBLISHERS:

University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania – Faculty of Biotechnologies

Address: 59 Mărăști Blvd., District 1, Zip code 011464, Bucharest, Romania,

Phone: + 40 21 318 25 64, Fax: +40 21 318 28 88,

E-mail: florentina.matei@biotechnologii.usamv.ro, Webpage: http://biotechnologyjournal.usamv.ro

CERES Publishing House

Address: 1 Piața Presei Libere, District I, Zip code 013701, Bucharest, Romania Phone: + 40 21 317 90 23, E-mail: edituraceres@yahoo.com, Webpage: www.editura-ceres.ro

Copyright 2016

To be cited: Scientific Bulletin Series F "Biotechnologies", Volume XX, 2016

The publisher is not responsible for the opinions published in the Volume. They represent the authors' point of view.

ISSN 2285-1364, CD-ROM ISSN 2285-5521, ISSN Online 2285-1372, ISSN-L 2285-1364

International Database Indexing: COPERNICUS, CABI, GOOGLE SCHOLAR, DOAJ, Scipio, PBN (Polish Scholarly Bibliography), OCLC, Research Bible, Cite Factor (Academic Scientific Journals), Universal Impact Factor

SUMMARY

AGRICULTURAL BIOTECHNOLOGY

| THE PARASITE BROOMRAPE (OROBANCHE CUMANA) IN SUNFLOWER – IDENTIFYING |
|---|
| SOURCES FOR GENETIC RESISTANCE- Gabriel Florin ANTON, Maria JOITA- |
| PACUREANU, Călina Petruța CORNEA |
| THE INFLUENCE OF THE FEMALE PARENT ON THE INDUCING RATE WITH FIVE |
| DIFFERENT INDUCER LINES IN MAIZE DH TECHNOLOGY - Ana Raluca BIȚICĂ, Aure |
| GIURA, Petruța CORNEA |
| PRELIMINARY RESULTS OF THE WINTER PEAS BREEDING PROGRAM - Ancuta |
| CRÎNGAŞU (BĂRBIERU) |
| GENETIC CONTROL OF GRAIN SIZE AND WEIGHT IN WHEAT - WHERE ARE WE NOW? |
| - Daniel CRISTINA, Matilda CIUCA, Petruta Calina CORNEA |
| THE INFLUENCE OF SEVERAL ABIOTIC FACTORS ON FUSARIUM SPP. BIOLOGY · |
| Roxana DUDOIU, Stelica CRISTEA, Daria POPA, Carmen LUPU, Maria OPREA |
| LEATHER HYDOLYSATE EVALUATED AS BIOACTIVE POTATO FERTILIZER - Mioara |
| Ancuța DUMITRU, Gabriel CORBU, Ștefana JURCOANE |
| INCREASING THE ANTIOXIDANT ACTIVITY, TOTAL PHENOLIC AND ASSIMILATORY |
| PIGMENTS CONTENT BY OPTIMIZING THE IN VITRO GROWTH CONDITIONS OF |
| LYCIUM BARBARUM PLANT- Mihaela DUŢU, Aurel ARDELEAN, Mirela ARDELEAN |
| Dorina CACHIŢĂ-COSMA, Marian BURDUCE, Andrei LOBIUC, Elida ROSENHECH |
| PRELIMINARY RESULTS REGARDING THE TESTING OF TREATMENTS WITH LIGHT |
| EMITTING DIODE (LED) ON THE SEED GERMINATION OF ARTEMISIA DRACUNCULUS |
| L Irina-Maria ENACHE, Oana LIVADARIU |
| AMOUNT OF DNA EXTRACTED FROM DIFFERENT TISSUES OF APPLE TREES IN |
| SPRINGTIME CAN BE USED TO DESCRIBE THE LEVEL OF INFECTION- Desareda |
| MERO, Ariola BACU |
| COMPARISON OF THREE DETECTION METHODS OF PHYTOPLASMA AT APPLE TREES |
| PROVES THE ADVANTAGE OF AMPLIFICATION OF SPECIFIC 16SrADN -Desareda |
| MERO, Ariola BACU, Margarita HYSKO |
| GETTING PLANTS NICOTIANA TABACUM THAT SIMULTANEOUSLY EXPRESS |
| HETEROLOGOUS GENE OF TWO ACYL-LIPID DESATURASES CYANOBACTERIUM |
| DESC AND DESA- Tetyana KYRPA-NESMIIAN |
| THE INFLUENCE OF GENOTYPE AND SOIL TILLAGE SYSTEM ON FORMING |
| PRODUCTION COMPONENTS FOR SWEET CORN UNDER SOIL-CLIMATE CONDITIONS |
| SPECIFIC TO ROMANIAN PLAIN - Maria Cristiana ONCICĂ (MORARU), Ricuța-Vasilica |
| DOBRINOIU, Luminița VISAN, Silvana DANAILA (GUIDEA), Izabela OPREA |
| THE INFLUENCE OF TECHNOLOGICAL LINKS UPON SWEET CORN YIELD QUALITY |
| UNDER SOIL-CLIMATE CONDITIONS OF THE ROMANIAN PLAIN - Maria Cristiana |
| ONCICĂ (MORARU), Ricuța-Vasilica DOBRINOIU, Luminița VISAN, Silvana DANAILA- |
| GUIDEA, Andrei Gabriel IVAN |
| GENETIC ANALYSIS OF QTLs CONTROLLING CARPEL NUMBER IN CITRUS - Osman |
| GULSEN, Aydin UZUN, Ihsan CANAN |
| PHYTOHORMONE-LIKE PRODUCING BACILLUS INCREASE TOMATO SEEDLINGS |
| QUALITY - Oana-Alina SICUIA, Sorina DINU, Florica CONSTANTINESCU |

| INFLUENCE OF DIETHANOLAMINE SALT OF 4-NITROBENZOIC ACID IN CALLUS | |
|---|-----|
| CULTURE AT MOMORDICA CHARANTIA L Alina SIMINA, Manuela CRISAN, Sorin | |
| CIULCA, Dorica BOTAU | 89 |
| EXTRACTION AND ANALYTICAL METHODS OF CAPSAICINOIDS - A REVIEW - Roxana- | |
| Mădălina STOICA, Mișu MOSCOVICI, Caterina TOMULESCU, Narcisa BĂBEANU | 93 |
| ACCELERATED MICROPROPAGATION OF ENDEMIC FRITILLARIA AUREA SCHOTT - | |
| Suleyman KIZIL, Tahsin SOGUT, Ugur SESIZ, Khalid Mahmood KHAWAR | 99 |
| INFLUENCE OF DIFFERENT IN VITRO SIMULATORS FOR HYDRIC STRESS FOR | |
| GROWTH AND DEVELOPMENT OF POTATO - Andreea TICAN, Mihaela CIOLOCA, | |
| Nicoleta CHIRU, Carmen Bădărău | 105 |

FOOD BIOTECHNOLOGY

| EFFECTS OF TREATMENTS WITH ASCORBIC ACID AND GLUTATHIONE ON THE | |
|---|-----|
| SAUVIGNON BLANC WINE COLOUR DURING BOTTLE AGING - Gianina Antonela | |
| BADEA, Arina Oana ANTOCE | 113 |
| COLOR CHANGES DURING THE STORAGE IN BOTTLES OF MUSCAT OTTONEL WINE | |
| TREATED WITH ASCORBIC ACID AND GLUTATHIONE - Gianina Antonela BADEA, | |
| Arina Oana ANTOCE | 121 |
| PHYSICAL CHEMICAL STUDIES REGARDING CIDER STABILITY STORED UNDER | |
| AMBIENT CONDITIONS - Iuliana BELEUZU BODA, Marius Cristian BODA | 129 |
| STUDY CONCERNING THE USE OF GERMINATED OAT FLOUR IN OBTAINING | |
| DIETARY BREAD - Dorica BOTAU, Panfil PIRVULESCU, Ersilia ALEXA | 137 |
| SOME PROPERTIES OF CAROB POD AND ITS USE IN DIFFERENT AREAS INCLUDING | |
| FOOD TECHNOLOGY - Selda BULCA | 142 |
| MUSHROOM MYCELIA CULTIVATION ON DIFFERENT AGRICULTURAL WASTE | |
| SUBSTRATES - Mihai Bogdan NICOLCIOIU, Gabriela POPA, Florentina MATEI | 148 |
| PHYSICAL AND CHEMICAL CHARACTERISTICS OF CROSSED OLIVES AND THEIR | |
| CONVENIENCE TO GREEN TABLE OLIVE FERMENTATION BY USING Lactobacillus | |
| plantarum AS A STARTER CULTURE - Yasin OZDEMIR, Sefik KURULTAY | 154 |
| DETECTION OF GENETIC VARIABILITY IN Pleurotus eryngii USING TUBULIN-BASED | |
| POLYMORPHISM MOLECULAR MARKERS - Gabriela POPA, Catalina VOAIDES, Matilda | |
| CIUCA, CalinaPetruta CORNEA | 162 |
| SCREENING FOR S-LAYER PRODUCTION BY SOME LACTOBACILLI FROM HOME- | |
| MADE FERMENTED FOODS - Iulia-Roxana ȘTEFAN, Silvia-Simona GROSU-TUDOR, | |
| Medana ZAMFIR, Petruța-Călina CORNEA | 167 |
| EFFECT OF EXTRACTION VARIABLES ON THE OMEGA-3 EICOSAPENTAENOIC ACID | |
| (EPA) CONTENT OF (Nannochloropsis oculata) MICROALGA OIL - Osman Kadir TOPUZ, | |
| Adem KAYA, Ali Can ALP | 172 |
| ALGAL OIL: A NOVEL SOURCE OF OMEGA-3 FATTY ACIDS FOR HUMAN NUTRITION - | |
| Osman Kadir TOPUZ | 178 |
| STUDIES ON THE CHROMATIC CHARACTERISTICS OF SOME ROMANIAN RED WINES - | |
| Luminița Valerica VIȘAN, Ricuța Vasilica DOBRINOIU, Silvana DANAILA-GUIDEA, | |
| Diana GROPOSILA-CONSTANTINESCU | 184 |

FOOD SAFETY

| THE EVALUATION OF TRANSFER LEVEL FOR PESTICIDE RESIDUES, IN CASE OF | |
|--|-----|
| PROCESSING BY SOLVENT EXTRACTION OF MEDICINAL PLANTS - Veronica | |
| DRUMEA, Roxana NITA, Lenuta ZGLIMBEA, Laura OLARIU | 195 |
| THE INFLUENCE OF PHYSICAL PARAMETERS OF MALT IN THE QUALITY OF WORT | |
| FOR BEER - TanjaKAMBURI, Luljeta XHANGOLLI | 200 |
| INVESTIGATION OF SOME EXTRACTION METHODS FOR THE RECOVERY OF PEANUT | |
| PROTEINS FROM OILS AND FATS- Elena MIHAI (DRĂGHICI), Maria PELE | 204 |
| VEGETABLE OIL CONVERSION INTO CORE-SHELL BIOPRODUCTS FOR STORED | |
| GRAIN PROTECTION-Mariana POPESCU, Florin OANCEA, Mălina DEȘLIU-AVRAM | 210 |
| ESTIMATION OF QUALITY OF 4 MONOFLORAL HONEY SAMPLES: ACACIA HONEY, | |
| LIME HONEY, OILSEED RAPE HONEY AND RASPBERRY HONEY - Ramona Iuliana | |
| SZASZ-ZIMA, Monica ENACHE | 214 |
| PHYSICO-CHEMICAL AND MICROBIOLOGICAL CHARACTERIZATION OF WHEAT | |
| FLOURS ON ROMANIAN MARKET IN RELATION TO THE SHELF LIFE - Radiana | |
| TAMBA-BEREHOIU, Ciprian–Nicolae POPA, VasilicaSIMION, Rodica CULEA | 218 |
| IDENTIFICATION OF THE MOST RELEVANT QUALITY PARAMETERS FOR BERRIES - A | |
| REVIEW - Elisabeta Elena TĂNASE, Vlad Ioan POPA, Mona Elena POPA, Mihaela GEICU- | |
| CRISTEA, Paul POPESCU, MihaelaDRĂGHICI, Amalia Carmen MITELUȚ | 222 |

INDUSTRIAL AND ENVIRONMENTAL BIOTECHNOLOGY

| INDOOR CULTIVATION OF SELECTED OIL - CONTAINING CONSORTIA OF | |
|---|-----|
| PHOTOSYNTHETIC MICROORGANISMS FOR FURTHER BIODIESEL PRODUCTION; | |
| PRELIMINARY FINANCIAL EVALUATION - Ioan I. ARDELEAN, Damian MANEA | 237 |
| HPTLC PHENOLIC COMPOUNDS FINGERPRINT AND ANTIOXIDANT ACTIVITY OF | |
| SAMBUCUS EBULUS LEAVES AND FRUIT - Corina BUBUEANU, Ramona-Daniela | |
| PĂVĂLOIU, Fawzia SHA'AT, Angela CĂȘĂRICĂ | 243 |
| IN VITRO BIODEGRADATION OF KERATINIZED SUBSTRATES BY KERATINOPHILIC | |
| FUNGI - Mariana CĂLIN, Olguța DRĂCEA, Iuliana RĂUT, Gelu VASILESCU, Mihaela | |
| BADEA DONI, Melania Liliana ARSENE, Elvira ALEXANDRESCU, Diana | |
| CONSTANTINESCU-ARUXANDEI, Luiza JECU and Veronica LAZĂR | 248 |
| EVALUATION OF THE CAPACITY OF DIFFERENT MICROORGANISMS TO SOLUBILIZE | |
| SEVERAL COMPOUNDS OF PHOSPHOROUS AND ZINC - ANDREEA DOBRE, LAURA | |
| ALEXANDRA MARIN, CARMEN MANOLE, NICOLETA ANDREI, CĂLINA PETRUȚA | |
| CORNEA | 254 |
| OBTAINING GROWTH CURVES FOR Scheffersomyces stipitis STRAINS AND THEIR | |
| MODELING - Mustafa GERMEC, Fatma Kubra KARTAL, Hazal GULDALI, Merve | |
| BILGIC, Asli ISCI, Irfan TURHAN | 263 |
| FERMENTATION OF ACID-PRETREATED TEA PROCESSING WASTE FOR ETHANOL | |
| PRODUCTION USING Saccharomyces cerevisiae - Mustafa GERMEC, Irfan TURHAN, Ercan | |
| YATMAZ, Nedim TETIK, Mustafa KARHAN | 269 |
| SHORT TERM EXPERIMENTS ON SYNTHETIC WASTE WATER TREATMENT IN | |
| LABORATORY ACTIVATED SLUDGE SEQUENCING BATCH REACTOR - Mirela C. | |
| IORDAN and Ioan I. ARDELEAN | 275 |
| LIFE CYCLE ASSESSMENT (LCA) ON EUROPEAN SKIMMED MILK POWDER | |
| PROCESSING PRODUCTION PLANT - Magdalini KROKIDA, Maria TAXIARCHOU, | |
| Antonis POLITIS, Antonis PEPPAS, Konstantina KYRIAKOPOULOU | 280 |
| NITROGEN AND PHOSPHORUS REMOVAL FROM MUNICIPAL WASTEWATER USING | |
| CONSORTIA OF PHOTOSYNTHETIC MICROORGANISMS - Roxana Gh. MANEA and Ioan | |
| I. ARDELEAN | 286 |

| GASTROPROTECTIVE POTENTIAL OF FAGUS SYLVATICA LEAVES EXTRACTS ON | |
|---|-----|
| STRESS-INDUCED ULCER MODEL ON RATS - Lucia PIRVU, Ioana NICU, Stelian | |
| SCHIOPU, Dragomir COPREAN | 293 |
| OPTIMIZATION OF HYDROLASE ENZYME DOSAGE IN THE PROCESS OF CELLULOSE | |
| HYDROLYSIS OF REJECT PULP AND BIOETHANOL FERMENTATION BY MICROBIAL | |
| CONSORTIUM - RATU SAFITRI, ATHENA DINANTY, GUSTAN PARI, MOH. | |
| NURZAMAN | 300 |
| LEVAN - A MINI REVIEW - Caterina TOMULESCU, Roxana STOICA, Claudia | |
| SEVCENCO, Angela CĂȘĂRICĂ, Mișu MOSCOVICI, Adrian VAMANU | 309 |

BIOTECHNOLOGY IN VETERINARY MEDICINE

| POPP, Doina DANES | 329 |
|--|-----|
| Rodica GURAU, Stelian BARAITAREANU, Marius Andrei MANESCU, Mihaela Cristiana | |
| POLYMERASE CHAIN REACTION TECHNIQUE OF PAN-SIMBU VIRUS GROUP - Maria | |
| STUDIES CONCERNING THE OPTIMISATION OF REVERSE TRANSCRIPTION | |
| MANASRA, Stelian BARAITAREANU, Maria Rodica GURAU, Doina DANES | 321 |
| Mohammad FARRAJ, Tamer ESSAWI, Israr SABRI, Kamel ADWAN, Issa RUMI, Assad | |
| MILK SHEEP AND GOAT IN WEST BANK PART OF PALESTINE - Elena AWWAD, | |
| VALIDATION OF RT-QPCR TECHNIQUE FOR DETECTION OF BRUCELLA GENOME IN | |

MISCELLANEOUS

| MICROBIAL XYLANASE: A REVIEW - Aglaia BURLACU, Călina Petruța CORNEA, | |
|--|-----|
| Florentina ISRAEL-ROMING | 335 |
| NEW BIOACTIVE COMPOSITES BASED ON BACTERIAL CELLULOSE AND NATURAL | |
| PRODUCTS - Angela CASARICA, Corina BUBUEANU, Ana Despina IONESCU, Ioana | |
| NICU | 343 |
| SELECTIVE FRACTIONS WITH ANTIOXIDANT ACTIVITY FROM ROMANIAN | |
| CULTIVATED CYNARA SCOLYMUS L Svetlana COLCERU-MIHUL, Sultana NITA, Alice | |
| GRIGORE, Corina BUBUEANU, Elena DRAGHICI, Emanuel VAMANU, Domnica | |
| RUGHINIS | 347 |
| THE DYNAMICS OF LIVE AND DEAD CELLS, AND COLONY FORMING UNITS OF | |
| E.COLI DH5 α STRAIN KEPT FOR FIVE DAYS AT 18°C IN MICROCOSMS CONTAINING | |
| FILTERED AND UNFILTERED SEA WATER - Roxana Gabriela CRISTINA and Ioan I. | |
| ARDELEAN | 352 |
| SATISFACTION LEVEL OF STUDYING THROUGH E-LEARNING SYSTEM AT THE | |
| STUDENTS ENROLLED IN THE BIOTECHNOLOGY EDUCATION - Gabriela MARGARIT, | |
| RaduCristian TOMA, Diana GROPOSILA, Dana BARBA | 358 |
| ANTIMICROBIAL ACTIVITY OF TEXTILES TREATED WITH ROSEMARY AND ORANGE | |
| ESSENTIAL OILS AGAINST A SELECTION OF PATHOGENIC FUNGI - Ovidiu | |
| IORDACHE, Andreea COZEA, Elena VĂRZARU, Elena STOICA, Corneliu PLATON, | |
| Steliana RODINO, Iuliana DUMITRESCU | 362 |
| BENEFITS OF THE E-LEARNING PLATFORMS AND CLOUD COMPUTING IN THE | |
| BIOTECHNOLOGY EDUCATION - Radu Cristian TOMA, Gabriela MARGARIT, Diana | |
| GROPOSILA, Dana BARBA | 370 |
| CULTIVATION OF FLORIDA OYSTER MUSHROOM ON VARIOUS TYPES OF | |
| SUBSTRATE - Denisa STĂNESCU, Emanuel VAMANU | 374 |