STUDIES REGARDING THE TRANSMISSION OF POTATO VIRUS Y (PVY) TROUGH SEVERAL MECHANICAL MEANS

BĂDĂRĂU C.L.^{1,2)}, STEFAN M.¹⁾, BĂRĂSCU N.¹⁾

¹⁾ National Institute of Research and Development for Potato and Sugar Beet Brasov ²⁾ Faculty of Food and Tourism, Transilvania University, Braşov, Romania

Keywords: PVY strains, PVY^N, PVY^{NTN}, mechanical transmission

ABSTRACT

The aim of this studies was to investigate the transmission of PVY strains PVY^o, PVY^N, PVY^{NTN}, via tuber cutting and plant treated in susceptible cultivars. For the tuber cutting experiment, after one infected tuber was cut with a knife, four uninfected tubers were cut sequentially with the same instrument without disinfecting it between the cuts. In the other experiments, the virus transmission from infected to healthy plants was made by bouncing, brushing, hammering, squeezing and carborundum rubbing treatments. These treatments allowed exchange of sap between the healthy and infected material. Results revelead that seed cutting did not transmit the pathogen, whereas the other plant treatments caused varying level of PVY transmission, depending on the experiments variant. Plant bouncing was the least effective whereas hammering was the most effective variant.

Abbreviations: PVY potato virus Y

INTRODUCTION

Potato virus Y (PVY), genus Potyvirus, member of the Potyviridae family is a growing concern for potato crops across many areas of the world [3,13,15,16,17,19]. The increase in PVY levels and the emergence of different strains of PVY, including the tobacco veinal strain PVY^N, the recombinant N:O strain PVY^{N:O} and the potato necrosis strain PVY^{NTN} have significant impact on both seed and commercial potato productions [6,7,10,11]. In recent years, some of the growers in the world observed unacceptable levels of current season spread of PVY, depending upon management practices [5,9]. PVY is transmited chiefly within the field by the movement and probing aphids. The significance of aphid-mediated current season spread of PVY has been well studied and documented [12,14]. However, information on current season spread of PVY through mechanical means is very limited. Recently, growers started asking questions about mechanical transmission of PVY because of guicker spread of some of the strains of PVY. Mechanical transmission may occur due seed-cutting and/or plant/tuber wounding in the standing crop. Sturz et al. (2000) reported that there was no mechanical transmission of PVY^o following the cutting of the infected tubers by hand or mechanical cutting [18]. In contrast, there are other reports suggesting mechanical transmission of PVY due to plant wounding and seed cutting [4]. Moreover, these observations were conducted mainly on PVY^O strains, wich has been replaced by PVY^{N:O} and/or PVY^{NTN} in some potato production areas. Farm operations such as post-emergence tillage and frequent use of tractors for spraying of the pesticidesand mineral oils during the crop season may cause plant wounding and sap exchange between the healthy and the infected plants. In adition, bouncing of the infected plants against the healthy ones or vice versa by strong winds may cause plant wounding, wich may potentially lead to virus transmission.

The study aimed to investigate the transmission of several PVY strains due to tuber cutting and to the plants treatments (on behalf to sap exchange between the healthy and infected material by bouncing, metal brishing, hammering, carborundum rubbing and squeezing). Our results demonstrated that PVY does not transmit via tuber cutting but it does via plant touch in certain situations.

MATERIAL AND METHOD

Plants material. Solanum tuberosum L. plantlets of varieties Hermes and Carrera were grown at the Biotechnology Department of National Institute of Research and Development for Potato and Sugar Beet Brasov, in summer of 2015 and tested for PVY strains. Plants tested positive were marked and tubers were harvested from individual plants and maintained separately. This material was used as source of the inoculum for mechanical PVY mean transmission.

Transmission trough seed tubers cutting

The experiment was carried out under greenhouse conditions. 45 tubers were used (3 infected tubers each of PYY^O, PYY^N and PVY^{NTN}, and 36 virus-free tubers, each of varieties Hermes and Carrera). After one tuber infected with a specific strain was cross-cut with a knife, four virus-free tubers from the same variety were cut sequentially with the same knife without using a disinfection solution. Both pieces from each tuber were planted separately in pots (in 2 rows of 5 plants). After every two rows, a row of five PVY-free tubers was planted in the pots and used as negative control. So, 30 tubers including 15 controls were studied for each PVY strain. There were 3 strains, 90 tubers were tested for each cultivar. The plants obtained from this biological material were tested using DAS ELISA, 21 and 42 days after planting.

Transmission through plant touching

Mechanical transmission of 3 PVY strains (PVY^O, PYY^N and PVY^{NTN}), due to plant touching was investigated. PVY-free tubers of cultivar Hermes were planted in pots and maintained in the greenhouse. After 3 weeks, these plants were tested again using DAS ELISA to make sure that all the material was PVY-free. These plants were then subjected to the following treatments:

- **Inclining and bouncing** healthy plants against PVY infected ones. The healthy and infected biological material were maintained in two separate rows. The healthy plants were bounced against the infected ones. Plants were bounced three times every week. These treatments had the intention to mimic the modality which could incline and bounce plants in the field.
- **Metal brushing** The leaf of an infected plant was brushed, followed by brushing on the leaf of the healthy plant; two leaves of the healthy plant were brushed. The brush had the sharp needles, which were supposed to exchange sap between the healthy and the infected leaflets.
- **Hammering** A leaf from an infected and a healthy plant was layered and pressed; 4 leaves from each healthy plant were hammered. Hammering could exchange sap between the healthy and the infected leaves.
- **Carborundum rubbing** The carborundum was put on the leaf of the healthy plant and then this leaf was rubbed by the leaf of the infected plant; 4 leaves of healthy plant were treated. Carborundum is an abrasive widely used for virus inoculations.
- **Squeezing** Twigs of the healthy plant and the infected were squeezed together. This may also exchange sap between the healthy and the infected leaves. Two twigs of the healthy plant were pressed using 2 rocks.

Therefore, 90 plants (3 strains x 6 treatments x 5 plants per treatment) were treated and tested.

Plants not treated were used as controls.

For each treatment, the touching procedure was repeated on a weekly basis three times. Two leaves from each plant were collected 7 days after the last touching and tested for PYY using DAS ELISA and PCR according to Lorenzen et al. (2006) [8]. In addition,

two daughter tubers from individual plants were also collected five weeks after the last touching treatment. Then, tubers from individual plants were composited, and tested for PYY using DAS ELISA to ensure translocation of virus from foliage to tubers. The experiment was repeated twice (experiment 1 starting in Mars 2015 and experiment 2 starting in Avril 2015).

DAS ELISA test. The plants were tested for the viruses using DAS-ELISA kits for PVY and according to the manufacturer's instructions with several exceptions (Bioreba, Switzerland). Also, the analysis was performed following the protocol described by [2] (100 μ l per well). Microplates were filled with substrate solution (p-nitro phenyl phosphate) incubated 1 hour for PVY and the absorbance values were estimated at 405 nm (A₄₀₅) on Tecan reader (Magellan softwere). The samples having A₄₀₅ values exceeding two times the average of healthy controls were considered virus infected. In the first stage, the material was tested for Potato virus Y (polyclonal antibodies) and then, this biological material was retested using monoclonal antibodies (mAb). The plates were coating with anti PVY-NOC mAb (Bioreba, Switzerland, antibodies that could recognize all the PVY strains excepting the PVY^O) and the virus was detected using alcalin phosphatase (AP) linked to anti–PVY-NOC mAb (Bioreba, Switzerland, specific for the strains PVY^N) [1].

RESULTS AND DISCUSSIONS

In the transmission through the seed tuber cutting experiment, none of the plants including daughter tubers of either Hermes and Carrera were positive except those used as sources of PVY inoculum (data not shown). This indicates that seed cutting is not effective in spreading PYY strains tested. These results are in close agreement with those of Sturz et al. (2000) [18], who reported no PYY^O transmission following the cutting of the infected tubers during seed multiplication. In contrast, Draper and Gudmestad (1992) [4] reported different degrees of transmission of PVY depending upon susceptibility of the cultivars and pattern of tuber cutting, such as infected eye to healthy eye, infected eye to healthy flesh, infected flesh to healthy flesh. They used cultivars Russet Norkotah, Shepody and Red LaSoda. Our study was conducted under greenhouse conditions, by using another cultivars and only one pattern of virus transmission (infected flesh to healthy flesh).

In the transmission through plant touching experiment, bouncing of healthy plants against the infected ones led to, out of five plants in each strain, two plants being infected with PYY^O and one plant with PVY^N in the first experiment, and one plant with PYY^O in the second experiment (Table 1). There was no transmission of PYY^{NTN} caused by bouncing. Out of five plants tested in each strain, metal brushing resulted in transmission of PVY^O in five plants in the first experiment and four in the second experiment, PVY^N in five plants in the first experiment and two in the second experiment and PYY^{NTN} in four plants in the first experiment and one plant in the second experiment. Hammering showed mechanical transmission of PVY in all the five plants tested in each strain in the first experiment and for PVY^N and PVY^{NTN} transmission, resulted four plants in the second experiment. Carborundum rubbing resulted in transmission of PVY in all the five plants tested in each strain in both experiments. Squeezing twigs of the healthy and infected plants together resulted in transmission of PVY^{O} in all five plants in the first experiment, and in four plants in the second experiment, PVY^{N} in two plants in the first experiment and four plants in the second experiment, and PVY^{NTN} in four plants in the first experiment and three plants in the second experiment. These results were also confirmed with RT-PCR (results not reported). The daughter tubers collected from positive plants (based on foliage testing) were also tested positive (data not shown), which confirmed that PVY spreads mechanically through different kinds of plant touching. The transmission of PVY strains due to hammering was identical to Carborundum rubbing in the first experiment and close

to Carborundum rubbing in the second experiment. Carborundum is most widely used for artificial disease inoculation. Thus, like Carborundum rubbing, hammering can also be used for artificial inoculation of PVY for experimental purposes.

Mechanical transmission of PVY strains trough plant touching treatments						
		Experiment 1*		Experiment 2*		
Treatments (mechanical means)	PVY strains	Number of plants tested	Number infected plants	Number of plants tested	Number infected plants	
Bouncing healthy plants against PVY infected	PVY ^O	5	2	5	1	
	PVY ^N	5	0	5	0	
	PVY ^{NTN}	5	0	5	0	
Metal brushing PVY infected plants and then healthy ones	PVY ^O	5	5	5	4	
	PVY ^N	5	5	5	4	
	PVY ^{NTN}	5	4	5	2	
Hammering leaf of the healthy and infected plant together	PVY ⁰	5	5	5	5	
	PVY ^N	5	5	5	4	
	PVY ^{NTN}	5	5	5	4	
Carborundum dusting on the healthy leaf plants + rubbing this with sap from infected leaf plant	PVY ^O	5	5	5	5	
	PVY ^N	5	5	5	5	
	PVY ^{NTN}	5	5	5	5	
Squeezing together twigs of healthy and infected plants	PVY ⁰	5	5	5	4	
	PVY ^N	5	2	5	4	
	PVY ^{NTN}	5	4	5	3	
Controls (untreated healthy plants)	PVY ^O	5	0	5	0	
	PVY ^N	5	0	5	0	
	PVY ^{NTN}	5	0	5	0	
*Experiment 1	was started in M	larch 2015 and F		started in Avril	2015	

Mechanical transmission of PVY strains trough plant touching treatments

Table 1

*Experiment 1 was started in March 2015 and Experiment 2 started in Avril 2015

The touching treatments used in this study would like to mimic field operations such as cultivation and spraying.

Usually, the pesticides and mineral oils are frequently sprayed in potato crops for the management of different pests. The equipment used for these practices may damage twigs and outer cells of infected plants and carry sap with them, introducing sap into the healthy plants. Thus, early rouging of PVY infected plants becomes important to minimize the risk of mechanical as well as insect mediated PVY transmission. PVY might be transmitted mechanically through wounds due to plant-to-plant contact from wind or by farm equipment operation. In the mechanical transmission through plant touching, the virus may move along with the sap. It may not happen that quickly in the mechanical transmission through tuber-cutting. In the same time, foliage usually shows higher viral titer than the tubers. Maybe, this is the reason because seed-cutting could not transmit PVY but plant touching did transmit it. Indeed, further research is required to know the mechanism of PVY transmission via plant touching.

CONCLUSIONS

The study showed that the seed cutting did not transmit the pathogen, whereas the other plant treatments caused varying level of PVY transmission, depending on the experiments variant. Thusly, different plant touching treatments (means that could mimic the field operations) induced mechanical transmission of PVY strains.

Regarding the efficiency of the plant treatment variants, the plant bouncing was the least effective whereas hammering was the most effective mechanical mean.

However, the study should be repeated under field conditions to confirm the results.

Acknowledgements This work was supported by a grant of the Romanian National Authority for Scientific Research, CNDI-UEFISCDI, PN-II-PT-PCCA-2013-4-0452, project number 178/2014.

BIBLIOGRAPHY

1. **Bădărău, C.L., Damșa ,F., Olteanu, Gh., Mărculescu, A.,** 2015 - *Total ascorbic acid content in 10 varieties of potato different resistant to PVY necrotic strains.* In: *Challenges in medicine, food control and environmental.* Floroian L. and Badea M. (ed.), Transilvania University from Braşov Publisher, p. 191-192.

2. Clark, M.F., Adams, A.N., 1977 - Characterization of the microplate method of the enzyme-linked immunosorbent assay for the detection of plant virus. Journal of General Virology 34: 475-483.

3. Davidson, R.D., Houser, A.J., Sather, K., Haslar, R., 2013 - Controlling PVY in seed: what works and what does not. American Journal of Potato Research 90: 28-32.

4. **Draper, M.A., and Gudmestad, N.C.,** 1992 - *Mechanical transmission of PVY by seed cutting,* American Potato Journal 69: 577-578.

5. Fageria, M.S., Singh, M., Nanayakkara, U., Pelletier, Y., Nie, X., Wattie, D., 2013 - *Monitoring current season spread of potato virus* Y *in potato fields using ELISA and realtime RT-PCR*, Plant Disease 97: 641-644.

6. Gray, S., De Boer, S., Lorenzen, I., Karasev, A., Whitworth, J., Nolte, P., Singh, Boucher, R.A., Xu, H., 2010 - *Potato virus* Y : an evolving concern for potato crops in the United States and Canada. Plant Disease 94: 1384-1397.

7. Hu, X., He, C. Xiao, Y., Xiong, X., Nie, X., 2009 - Molecular characterization and detection of recombinant isolates of potato virus Y from China. *Archives of Virology* 154: 1303-1312.

8. Lorenzen, J.H., Piche, L., Gudmestad, N.C., Meacham, T.P.H., Shiel, P.J., 2006 - A *multiplex PCR assay to characterize potato virus Yisolates and identify strain mixtures*. Plant Disease 90: 935-940.

9. MacKenzie, T.D.B., Fageria, M.S., Nie, X., Singh, M.. 2014 - Effects of crop management practices on current season spread of potato virus Y. Plant Disease 98: 213-222.

10. Nie, B., Singh, M., Murphy, A.M., Sullivan, A., Xie, C., Nie, X., 2012 - Response of potato cultivars to five isolates belonging to four strains of Potato virus Y, Plant Disease 96: 1422-1429.

11. Nie, X., Singh, M., Pelletier, Y., McLaren, D.L., 2013 - *Recent advances on potato virus Y research in Canada.* American Journal of Potato Research 90: 14-20.

12. **Pelletier, Y., Nie, X. Giguere, M.-A., Nanayakkara, U.N., Maw, H.E.L., Foottit, R.G.,** 2012 - A New Approach for the identification of aphid vectors (Hemiptera: Aphididae) of Potato virus Y. Journal of Economic Entomology 105: 1909-1914.

13. **Quenoille, J., Vassilakos, N., Moury, B.,** 2013 - Potato virus Y: a major crop pathogen that has provided major insights into the evolution of viral pathogenicity, Molecular Plant Pathology, 14: 439.

14. Radcliffe, E.B., and Ragsdale, D.W., 2002 - *Invited review. Aphid transmitted potato viruses: the importance of understanding vector biology.* American Journal of Potato Research 79: 353-386.

15. Rasocha, V., Hausvater, E., Dolezal, P., 2008 - *Harmful agents of Potato. Abionoses, Diseases, Pests.* Potato Research Institute Havlickuv Brod Ltd (ed.), pp. 66-67.

16. Scholthof, K.B., Adkins, S., Czosnek, H., Palukaitis, P., Jacquot, E., Hohn, B., Saunders, K., Candresse, T., Ahlquist, P., Hemenway, C., Foster, G.D., 2011 - *Top 10 plant viruses in molecular plant pathology*, Molecular Plant Pathology, 12: 938-954.

17. Steward, G., De Boer, S., Lorenzen, K., Withworth, J., Nolte, P., Singh, R., Boucher, A., Xu, H., 2010 - Potato virus Y an evolving concern for potato crops in the United State and Canada, Plant Disease, 94(12): 1384-1397.

18. **Sturz, A.Y., Stewart, J.G., McRae, K.B., Diamond, J.F., Lu, X., Singh, R.P.,** 2000-. Assessment of the importance of seed cutting, inseason cultivation, and the passage of row equipment in the spread of PVYO in potatoes. Canadian Journal of Plant Pathology 22: 166-173

19. Wrobel, S., Wlasik, A., 2013 - Seed Potato production in Poland, American Journal for Potato Research, 3: 260-268.

UNIVERSITY OF CRAIOVA FACULTY OF AGRONOMY in collaboration with UNIVERSITY OF BELGRADE FACULTY OF AGRICULTURE

in collaboration with

Romanian Academy of Agriculture and Forestry - Craiova Branch



SCIENTIFIC INTERNATIONAL CONFERENCES

The 12th ANNUAL MEETING "DURABLE AGRICULTURE – AGRICULTURE OF THE FUTURE"

PARTICULAR FOCUS OF THE CONFERENCE

"ADVANCED METHODS FOR A SUSTAINABLE AGRICULTURE, SILVICULTURE AND FOOD SCIENCE"

VOL. XLVI/1/2016

ISSN 1841-8317 ISSN CD-ROM 2066-950X

17th-18th NOVEMBER 2016

CRAIOVA ROMANIA

Editorial Board

Professor Cezar Ionuț SPÎNU, PhD., Rector of University of Craiova

Professor Aurel CĂLINA, PhD., Vice-Rector, University of Craiova, Faculty of Agronomy Professor Tudor ALEXANDRU, PhD., University of Craiova, Dean of the Faculty of Agronomy Professor associate Mariana NICULESCU, PhD., University of Craiova, Vice- Dean of the Faculty of Agronomy

Professor associate Sorin Petruţ BORUZ, PhD., University of Craiova, Vice-Dean of the Faculty of Agronomy

Professor associate Mugurel COLĂ, PhD., University of Craiova, Vice-Dean of the Faculty of Agronomy

Professor asociate, Liviu AureL OLARU, PhD., University of Craiova, Faculty of Agronomy

Editorial Review Board

Professor Aurel CĂLINA, PhD., Vice-Rector, University of Craiova Faculty of Agronomy Professor Tudor ALEXANDRU, PhD., University of Craiova, Dean of the Faculty of Agronomy Professor associate Mariana NICULESCU, PhD., University of Craiova, Vice- Dean of the Faculty of Agronomy

Professor Multiple Dr. H. C. Stefano GREGO, President of ESNA, World Agricultural Heritage Foundation - Chairman of Scientific Committee, Deputy Director of CIRPS, Italy

Professor asst. dr. M. Cuneyt BAGDATLI, University o Nevsehir Haci Bektas Veli Engineering and Architecture Faculty, Department of Biosystem Engineering, Turkey

Professor Vlado LICINA, PhD, Faculty of Agriculture, University of Belgrade, Vice-President of ESNA

Professor associate Sorin Petruţ BORUZ, PhD., University of Craiova, Vice-Dean of the Faculty of Agronomy

Professor associate Mugurel COLĂ, PhD., University of Craiova, Vice-Dean of the Faculty of Agronomy

Professor asociate, PhD. Liviu AureL OLARU, University of Craiova, Faculty of Agronomy Professor Michael PÖSCHL, PhD., University of Agriculture and Forestry in Brno, Czech Republic

Professor Ivan ILIEV, PhD., University of Forestry, Sofia, Bulgaria

Professor Christos KARELAKIS, Democritus University of ThraceSchool of Agricultural and Forestry Sciences, Greece

Professor associate LIDIA MISHEVA, PhD.- Institute of Soil Science, Sofia, Bulgaria Professor Gheorghe SIN, PhD., President of the Agricultural and Forestry Sciences Academy "Gheorghe Ionescu Şişeşti", Romania

Professor Mihai NICOLESCU, PhD., Vice-President of the Agricultural and Forestry Sciences Academy "Gheorghe Ionescu Şişeşti"

Professor Mircea MIHALACHE, PhD., U.S.A.M.V. Bucureşti, Faculty of Agriculture Professor associate Jenica CĂLINA, PhD., Faculty of Agronomy, University of Craiova Professor Cristina BĂBEANU, PhD., University of Craiova, Faculty of exact Sciences

Professor Rodica BERCU, PhD., "Ovidius" University of Constanţa, Faculty of Natural Sciences and Agricultural Sciences

Professor, Markovic NEBOJSA, PhD., Faculty of Agriculture, University of Belgrade, Serbia Professor associate Dumitru ILIE, PhD., University of Craiova, Faculty of Mechanical Engineering Professor associate IMBREA FLORIN, PhD., Banat University of Agricultural Sciences and Veterinary Medicine "Regele Mihai I al României" of Timisoara, Romania

Professor associoate Marius FĂGĂRAŞ, PhD., "Ovidius" University of Constanţa, Faculty of Natural Sciences and Agricultural Sciences

Dr. Valentin VLÅDUŢ, INMA Bucureşti

Dr. Cristiana OPREA, Joint Institute of Nuclear Research, Dubna Moscow, Russian Federation Doc. Dr. Zoran PRŽIĆ ZORAN, University of Belgrad, Faculty of Agriculture, Vice-President of the E.S.N.A., Serbia

Professor associate Mihnea GLODEANU, PhD., Faculty of Agronomy, University of Craiova Professor associate Constantin NEŢOIU, PhD, Faculty of Agronomy, University of Craiova

Dr. Mirela RIDICHE, Oltenia Museum

Junior Lecturer Alexandru SĂRARU, PhD., Faculty of Agronomy, University of Craiova Prof. dr. Jörg RINKLEBE, University of Wuppertal, Commission Head of Water- and Waste-Management / Sanitary Environmental Engineering, Germany

Secretary

Junior Lecturer Marius MILUT, PhD., University of Craiova, Faculty of Agronomy Junior Lecturer Dragos MEDELETE, PhD., University of Craiova, Faculty of Agronomy Junior Lecturer Alin CROITORU, PhD., University of Craiova, Faculty of Agronomy Junior Lecturer Marius CIOBOATĂ, PhD., University of Craiova, Faculty of Agronomy Junior Lecturer Andi CIOBANU, PhD., University of Craiova, Faculty of Agronomy

CONTENTS - VOL. XLVI/1/2016

Working Group No. 1

PLANT CULTIVATION AND ANIMAL GROWING TECHNOLOGIES

BĂDĂRĂU C.L., RAKOSY-TICAN E., AURORI A., CHIRU S.C., OLTEANU GH., STEFAN M.,
GHINEA A The incidence of potato virus y (necrotic strains) in seed potato grown in several
Romanian Counties (preliminary studies) 7 BĂDĂRĂU C.L., STEFAN M., BĂRĂSCU N Studies regarding the transmission of potato virus y
(PVY) trough several mechanical means
BARASCU NINA, ȘTEFAN FLORIANA MARIA, HERMEZIU RADU, BADARAU CARMEN LILIANA
 Study on the culinary and technological quality appreciation of new varieties of potato obtained at NIRDPSB Brasov 22
NIRDPSB Brasov 22 ZORAN BEŠLIĆ, SLAVICA TODIĆ, NEBOJŠA MARKOVIĆ, ZORAN PRŽIĆ - Influence of early
basal leaf removal on yeald components and must quality on CV. Sauvignon Blanc31BONCIU ELENA, SĂRAC IOAN - Implications of modern biotechnology in the food security and food31
36 BONCIU ELENA - Basic raw materials used in processing of the snack food (ecological/non
ecological) and their expanding capacity
BONEA DORINA - The effect of climatic conditions on the yield and quality of maize in the central part of Oltenia
48 BONEA DORINA - Effect of the aqueous extracts of Amoracia rusticana L. on the seed germination
and seedling growth of Zea mays L. under drought stress
DANIELA DOLORIS CICHI, COSTEA DORIN CONSTANTIN, GHEORGHIU NICOLAE - The cold
hardiness of some varieties of grapevine cultivated in the viticultural area Plenita (Southwestern
Romania)
CICHI MIHAI - Result of the association variety x rootstock on the growth and development of aerial
and underground systems in the species of apple and pear
at some tree varieties on the soils from the region of Oltenia
variety
FLORICA COLĂ, VIORICA URECHEAN MUGUREL COLĂ - Behavioural indicators in the nutrition- excretion process of the Holstein-Friesian catlle exploited at S.C.D.A. Simnic
COLĂ MUGUREL - Results regarding the behavioural indicators during the automatic milking of the
dairy cattle exploited at S.C.D.A. Şimnic Craiova
MARIJA ĆOŠIĆ, RUŽICA STRIČEVIĆ, NEVENKA DJUROVIĆ, LJILJANA PROKIĆ, MILENA MARJANOVIĆ, DJORDJE MORAVČEVIĆ - Impact of irrigation regime and application of kaolin on
the stomatal conductance and leaf water potential of pepper and tomato
CONSTANTINESCU, MARIANA NICULESCU - The influence fertilization on the quality production
of rye in the conditions of sandy soils from southern Oltenia
CV. 'Polka' as dependent on the type of aboveground growth
MILICA DIMA, MIHAELA CROITORU, RETA DRĂGHICI, EMILIA CONSTANTINESCU, MARIANA NICULESCU - Researches regarding on the efficacy and selectivity of herbicides to combat weeds
from lavender on sandy soils
DOBRE CL.M., COTIGĂ C Adapting perennial graminacee production for hills`area of Oltenia
region 119 DOBRE CL.M., COTIGĂ C Organic and mineral fertilization of temporary grasslands – ecological
measure and the efficiency on enhancing fodder production scenario in the hills`area of Oltenia
region
ŻELJKO DOLIJANOVIĆ, MILENA SIMIĆ, NEBOJŠA MOMIROVIĆ, ĐORĐE MORAVČEVIĆ,
BILJANA JANOŠEVIĆ - The effects of different cover crops on grain yield of popcorn (Zea mays L.
ssp. everta sturt) 129 DRĂGHICI IULIAN, RETA DRĂGHICI, MIHAELA CROITORU - Capitalizing of the sandy soils from
southern Oltenia through cultivation of new varieties of triticale

RETA DRAGHICI, AURELIA DIACONU, DRAGHICI IULIAN, MILICA DIMA, MIHAELA CROITORU, MARIETA PLOAE, COTEȚ GHEORGHE, ALINA PARASCHIV - Results on the influence space nutrition on growth and development of sweet potato plant under sandy soils in southern Oltenia
RETA DRAGHICI, DRAGHICI IULIAN, AURELIA DIACONU, MILICA DIMA - Variability of genetic
resources of cowpea (Vigna unguiculata) studied in the sandy soil conditions from Romania
ROXANA DUDOIU, VIOREL FATU, CARMEN LUPU, DARIA POPA, ELENA RADU, MARIANAPOPESCU - Antimicotic activity of Ocimum basilicum essential oil against stored fungi154
IANCU PAULA, SOARE MARIN, BONEA DORINA, PĂNIȚĂ OVIDIU - Preliminary results
concerning the genetic variability of some mutant lines of wheat experimented to ARDS Caracal 159 NICOLAE IONESCU, FLORIAN TRAȘCĂ, GINA MINCĂ, GEORGETA TRAȘCĂ, IULIA CIODARU AND MARIA VOICA - Fruit characteristics of Digitaria sanguinalis (L.) scop. weed from maize crop 169 TATJANA JOVANOVIĆ CVETKOVIĆ, DRAGUTIN MIJATOVIĆ, IVANA RADOJEVIĆ, ZORICA
RANKOVIĆ-VASIĆ, DRAGAN NIKOLIĆ, IVANA MOŠIĆ - The characteristics of fruitfulness of some
interspecies grapevine cultivars
parameters on uvological characteristics of variety 'Blatina'
KAKHA NADIRADZE, NANA PHIROSMANASHVILI - An importance of virus free seed production
systems in Georgia 18 CHRISTOS KARELAKIS, EFSTRATIOS LOIZOU, MARIUS VLADU - Exploring the
competitiveness, of the Greek fisheries sector 188 KRGA I., SIMIĆ A., BIJELIĆ Z., MANDIĆ V., VASILJEVIĆ S., KARAGIĆ Đ., MILIĆ D Interspecies
relations and yield of different field pea/oats mixtures
MARKOVIĆ, N., PRŽIĆ, Z., TODIĆ, S., BEŠLIĆ, Z Productive and technological characteristics of table varieties growe in the conditions of Oplenac vineyards
SAŠA MATIJAŠEVIĆ, ZORAN BEŠLIĆ, ZORAN PRŽIĆ, DRAGOLJUB ŽUNIĆ, SLAVICA
TODIĆ,NEBOJŠA MARKOVIĆ,ZORICA RANKOVIĆ-VASIĆ, BRATISLAV ĆIRKOVIĆ, VERA VUKOSAVLJEVIĆ, DUŠICA ĆIRKOVIĆ, MERSIJA DELIĆ - Influence of cultivar characteristics of
Muscat table grapevine cultivars (Vitis vinifera L.) on grape brandy composition and quality
UGRINOVIĆ, M Effects of continuous fertilization on the cucumber seedling quality
KLAJDI NALLBANI, TURMALA J., ENSTELA SHUKULLARI, REXHEP BAJRAMAJ - Hormonal
treatment to prevent the residue of placenta in cows 22: NIŢU SORINA, ȘTEFAN FLORIANA MARIA, CHELMEA CARMEN, HERMEZIU MANUELA - 22: Draliminaru studies en mainteining the biodiversity of madicinal plants within NIDDESE Preserver 22:
Preliminary studies on maintaining the biodiversity of medicinal plants within NIRDPSB Brasov 228 C.V. POPESCU, Claudia BORLEANU, C. BORA - Measurements concerning some yield elements 238 on sunflower cultivars in the climatic conditions of 2016 at S.C.D.A. Simnic 238
IVANA RADOJEVIĆ, MILOŠ RISTIĆ, MLAĐAN GARIĆ, TATJANA JOVANOVIĆ CVETKOVIĆ, ZORICA RANKOVIĆ-VASIĆ, DRAGAN NIKOLIĆ, ZORAN PRŽIĆ - Yield and quality of Müller-
Thurgau clone GM11 of Nis grape growing region 242 RĂDULESCU DIANA, TĂNASIE ȘTEFANIA ELIZA, DUMITRU FLORIN - Correlations between some physiological processes of maize hybrids with Minerva and Danubiu influence on yield and
quality of grain 240
RĂDULESCU DIANA, TĂNASIE ȘTEFANIA ELIZA, DUMITRU FLORIN - Differentiated application
of nitrogen and phosphorus cancause physiological traits in maize
agrobiological and technological characteristics of promising grapevine genotypes obtained for red
wine production
C.A. ROŞCULETE, ELENA ROŞCULETE - The influence of foliar fertilizers on rape crops from SCDA Caracal
ELENA ROȘCULETE, ROȘCULETE C. A The influence of freezing on the vitamin C content in some vegetable products
SĂLCEANU C., OLARU L Researches on weed control on strawberries
the lemon juice after sweetening task
development of organic farming systems in Romania as the basis for obtaining of innocuity
agricultural raw materials – used in functional food
the knowledge of the morphology, ecology and damage caused by mining moth chestnut (Cameraria
orhidella Deschka & Dimic) in the Husi - Vaslui areal conditions from eastern of Romania

TĂLMACIU NELA, TĂLMACIU MIHAI, MIROIU CARMEN, HEREA MONICA - Observations on the	
biology of the Cameraria ohridella Deschka Dimič species (leaf mining moth), dangerous pest of the	
chestnut tree	290
MARIUS VLADU, CRISTINA-EMANUELA VLADU - Studies on the implementation of the	
clasification system "SEUROP" to award quaity classes of carcasses of pig	297
Xhafa A., Margariti K., Kumbe I Seroprevalence of Toxoplasma gondii infection in pigs in	
extensive breeding from Saranda district of Albani	304
ZELJKOVIĆ SVJETLANA, ŠUŠAK UROŠ, TODOROVIĆ VIDA - Influence of two biostimulants	
application on growth and development of Tropaeolum majus L.	308

Working Group No. 2

SOIL SCIENCES

EUGENIA GAMENT, MARIANA MARINESCU, VERA CARABULEA, GEORGIANA PLOPEANU -	
Steel slag, a substitute of liming materials in agriculture	314
GRECU FLORINA, PAVEL STEFAN - Pedological and chemical characterization of a brown -	
stagnic vertosoil in order to establish the fertilization plan	322
ANČA-ROVENA LĂCĂTUȘU, RADU LĂCĂTUȘU, MIHAIL DUMITRU, IRINA-RAMONA MORARU, ANDREI VRÎNCEANU, CLAUDIA BĂLĂCEANU, LAVINIA BURTAN - Decontamination of a	
petroleum hydrocarbons polluted soil by different bioremediation strategies	326
MARIANA MARINESCU, ANCA LACATUSU, EUGENIA GAMENT, GEORGIANA PLOPEANU,	
VERA CARABULEA, MARINESCU MIHAI - A review of biological methods to remediate crude oil	
polluted soil	335
MATEI GABI-MIRELA, MATEI S., MOCANU VICTORIA, DUMITRU SORINA - Microbiological	
characterization of suppressive forest soil from Enisala	341
MATEI S., MATEI GABI-MIRELA, DUMITRU SORINA, IGNAT P Research on the role of microbial	
consortium in biosynthesis of humic precursors based on secondary exometabolites	348
POPESCU CRISTIAN - The natural condition of formation and the main features of soils from Farcas	
locality, district Dolj	357
POPESCU CRISTIAN - The cropping characterization of soils from Vladaia locality, district Mehedinti	364
RUJOI B., F. BARCA - Possibilities of soil regeneration using the caustobiolites in early stages of	
carbonization	372
CARMEN SIRBU, CIOROIANU TRAIAN, ADRIANA GRIGORE, ANA-MARIA STANESCU,	•••=
BOGDAN RUJOI, NICOLETA MARIN, LAVINIA BURTAN, MONICA DUMITRASCU, EMILIA NICU	
- Obtaining and testing of fertilizers with organic substances	376
TUDOSOIU C., ACHIM ELENA., GANEA-CHRISTU I Analytical calculation of water retention	570
capacity, specific for forestry terraces situated on slope degraded lands	382
	J0 Z
VUJOSEVIC A. - The influence of substrate composition on the development of seedlings of annual flavore Cetherarthy a reasonal	204
flowers Catharanthus roseus L.	391



