

BEHAVIOUR OF DIFFERENT POTATO VARIETIES BY SIMULATING IN VITRO OF HYDRIC STRESS CONDITIONS

Andreea TICAN¹, Mihaela CIOLOCA¹, Nicoleta CHIRU¹, Carmen BĂDĂRĂU²

e-mail: tican_andreea@yahoo.com

Abstract

The water deficit, extreme temperatures and low atmospheric humidity lead to drought, which is one of the limiting factors affecting crop quality and quantity. *In vitro* selection may shorten considerably the time selection of desirable traits and completes selection in the field. *In vitro* tissue culture can be used to determine the drought tolerance of the various varieties assuming that there is a correlation between plant cells *in vitro* and *in vivo*. For determining the resistance to water stress, the research was conducted in the Laboratory of Vegetal Tissue Culture and experience consisted of two factors: variety and nutritive medium with different osmotic agents. With the increasing amount of sorbitol and PEG, water absorption becomes difficult for plantlets from nutrient medium and thus was simulate the effect of drought over microplants. With the increasing amount of sorbitol or PEG into the growth medium, the drought intensity was bigger. Observations were made for different parameters: the average number of leaves; the average number of internodes; the average height of plantlet (cm); the average root length (cm); the average weight of fresh plantlet (mg); the average fresh root weight (mg). This study was performed in order to determine the best osmotic agent put into the *in vitro* medium for simulate the water stress, as well as for testing the particular genotypes genotypes on different culture media causing selection pressure of water stress.

Key words: water stress, *in vitro*, culture medium, PEG, sorbitol

Drought is one of the most important constraints for ecosystems productivity, in many regions around the world and water availability becomes more scarce for many agricultural areas. Among the factors influencing production can include insufficient rainfall, high levels of salts in the soil. Breeding varieties with drought tolerance is a challenge of great priority in the program of biotechnology. *In vitro* selection may shorten considerably time for selection of desirable traits and completes selection the field.

Drought can be defined in relation with rainfall deficit compared to potential evapotranspiration in a year, a specific season, or in relation with the exigencies of a particular culture at a certain moment. Also, drought may occurs in areas where annual and periodic rainfall are „normal” or more or less nearby to the multiannual average, but the water comes from rainfall rare with high intensities (mm / min) and do not permit its accumulation in the soil. This type of drought is more common in southern and eastern Romania with a tendency of extending in central and western country (Petcu, 2008).

Actions for a short period of stress induce reactions in plants of protection, adaptation (response) to respective conditions. In case of long action of stressogenic factors, reversible initial changes may get out of plant genome control and go into irreversible changes that lead to destruction of plant (Popovici A. *et al*, 2013).

Water deficit associated with temperatures over thermal threshold manifested in areas with climate temperate - continental leads several times to stagnating or even forced maturation (Morar, 1999).

There is some evidence that the roots are the primary sensors of water deficit in the soil, causing physiological and biochemical perturbations in stems; growth reduction can generally be interconnected with changes in plant nutrition.

Capacity of roots to pervade the soil depends on the power that roots can exert and may be associated with drought tolerance (Tardieu, 1994).

The effect of water stress on tuber formation depends on the stage of crop development. During stolonization phenophase, water stress is crucial. It reduces the number of tubers and total production,

¹ National Institute of Research and Development for Potato and Sugar Beet Brasov, Brasov, Romania

² Faculty of Food and Tourism, Transylvania University, Brasov, Romania

due to a decrease in stolons number (Usman, 2012).

Appearance of new approaches involving the use of high efficiency of several technologies (by applying several techniques: genetic, physiological, biochemical, molecular and biotechnology) offers hope for interesting innovations to maintain food security, poverty reduction and mitigating risk agriculture farmers in vulnerable environments (Obidiegwe *et al*, 2015).

In vitro selection may shorten considerably time for selection of desirable traits and selection completes in the field. For *in vitro* selection PEG, sucrose, mannitol and sorbitol were analyzed in several research papers, as agents of osmotic stress.

Taking into account the stress of drought on potato at NIRDPSB Brasov, Laboratory of Vegetal Cultures Tissue was conducted a study to explore tolerance / resistance to drought *in vitro* of five potato varieties.

With increasing amount of sorbitol and PEG, water absorption becomes difficult for plantlets from nutrient medium and thus simulates the effect of drought on microplants. With the higher amount of sorbitol or PEG into the growth medium, the drought intensity will be higher.

In vitro culture techniques minimizes external environmental changes due to nutrient medium defined and controlled conditions applying stress homogeneity.

MATERIAL AND METHOD

Microplants from the culture collection (belonging to five varieties) were multiplied to each internode and minicuttings were inoculated on Murashige-Skoog medium (1962) enriched with vitamins. After four weeks, the plants which were formed were multiplied to obtain nodal cuttings. Cuttings of these plants were used as explants for further multiplication *in vitro*. For rapid identification of genotype with resistance / tolerance to water stress was chosen two simulators of water stress. One of them was polyethylene glycol (PEG 6000) and the second was sorbitol (growth inhibitors). Both of them were separately added in medium using 4 concentrations.

Experimental conditions were that from specific growth chamber of plantlets provided in the working protocol; inoculation room of minicuttings is sterilized with laminar flow of sterile air; sterilization of the tubes is carried out in an oven at 180°C, and the culture medium sterilized must be inserted into test tubes in an autoclave at 120°C.

The results were examined by analysis of variance. Significance of differences was determined by method of multiple comparisons,

respectively Duncan test, stating the significance of any differences obtained by executing all comparisons possible, with the significant difference corresponding remoteness found in classification variants compared. Experimental differences higher than 5% are considered significant (Săulescu, N. A, Săulescu, N. N., 1967).

Experience included the following factors:

Experimental factor A: variety, with 5 graduations:

- a₁- Ruxandra;
- a₂- Sarmis;
- a₃- Gared;
- a₄- Marvis;
- a₅- Rustic.

Experimental factor B – nutrient media used (with different simulators of water stress) with 3 graduations:

- b₁-control medium MS, which contain was no osmotic agent;
- b₂- MS medium, to which was added PEG;
- b₃- MS medium, to which was added sorbitol.

Test tubes with minicuttings varieties proposed for *in vitro* testing of water stress (figure 1) were placed in the growth chamber, ensuring light and temperature regime needed growth and development of plantlets; after 4 weeks measurements were made for analysis of the following parameters: number of leaves and internodes/plantlet and plantlets height, root length and root fresh plant weight (mg).



Figure 1 Fresh inoculated minicuttings on MS medium to which was added on simulators of water stress

RESULTS AND DISCUSSIONS

Simulation drought *in vitro* was performed in order to identify the best varieties tolerant to drought. Observations revealed that was obtained a slowdown in regeneration cuttings.

By ample analysis of variety and nutritive medium (MS) with different osmotic agents on four concentrations (table 1), Duncan test indicated by multiple analysis, the different behavior for all variants tested.

Regarding the average number of leaves, the highest value was achieved by using control medium for Marvis variety (11 leaves) and by using sorbitol 1.5%, leaves obtained were on a

number of 11.25 (not differ statistically by control medium), but on maximum concentration of PEG 2% leaves were registered on a number of 10.5, which indicated that the variety has a strong tendency to resist on drought, by simulating the formation of a larger number of leaves. At the opposite pole was Sarmis variety, which for both inducers of water stress at concentrations of 2% number of leaves decreased dramatically (6 leaves). Water stress affected the absorption of nutrients from the nutrient medium, reduced photosynthesis process and had as effect decreasing the number of leaves.

The average number of internodes highlighted the Ruxandra variety on control medium, which recorded the highest number of internodes (6.75), but also was observed for Marvis variety tendency, to combat the effect of drought, which by application of sorbitol 1.5% in Murashige-Skoog nutrient medium, produced a greater number of internodes (6.50).

The average height of plantlet (cm) analyzed from the interaction of two factors - variety and nutrient medium with different concentrations of the two simulators of drought highlighted the use of control nutrient medium, with higher value for Sarmis variety of 13.625 cm (A), followed Marvis and Rustic varieties of 10.375 cm and 11.00 cm respectively.

In the case of water stress, plant root growth stops, water absorption can not be performed. Gared variety proved the root elongation, tendency to eliminate the effect of drought, produced by using sorbitol 1.5 and 1% (values of 10.750 cm and respectively 8.375 cm), exceeding even the medium option in which was not apply specific treatment for simulation drought (version control medium: 7.625 cm), indicating the combating drought. The same variety (Gared) by using PEG 1.5% for the induction of water stress, registered a high value for the average root length (7.125 cm), with no significant differences compared to root formed on control medium (7.625 cm). So, this variety resisted very well to water stress, by root elongation „ in search of water. "

The average weight of fresh plant, is contained in the general table of with interactions of variety and medium plus concentrations of drought simulators. The values were between 247.900 mg for Marvis variety (A) and 244.70 mg (A) for Ruxandra variety, both on control medium and 21.60 mg and 23.50 mg (values obtained for the same varieties Marvis and Ruxandra, but were used different osmotic agents, but with the same concentration (2%) PEG and sorbitol, indicating

that although Marvis and Ruxandra varieties obtained greater weight on control medium, PEG and sorbitol, at maximum concentrations had inhibitory effect on vigor plantlets and drastically reduced their weight (ex. even 10.4 times for variety Ruxandra).

The last analyzed element, fresh root weight (mg), had values between 163.625 mg for Ruxandra variety (A) on control medium and 123.625 mg by application sorbitol 0.5% (the same variety). Forming a vigorous root system, well branched in nutrient medium provides an access to nutrients substances.

Is was observed a drastic decrease in root weight, making them very fragile, using PEG at maximum concentration with values between from 4.1 mg (W) for Sarmis variety, 5.925 mg (W) for Marvis variety and 6,275 mg (W) for Rustic variety. That indicated that Sarmis, Marvis, Rustic varieties perceived on addition of PEG 2% as the most stressful factor for the growth of root for the weight values provided by multiple analysis by Duncan's test from A and B, and reached W and described the best effect of water stress for plantlets.




Figure 2 Effects of water stress stimulators (PEG and sorbitol) on average number of leaves and internodes in comparison with control medium

The process of forming leaves (figure 2) highlighted Marvis variety, both on control medium (with a number of 11 leaves), as well as media that had PEG (it were obtained 9.38 leaves) and sorbitol (8, 31 leaves).

Regarding the average number of internodes Rustic and Marvis varieties had the best resistance/tolerance by using sorbitol (forming 5.25 internodes / plant for both varieties).

Table 1

The combined influence of variety and inducer medium of *in vitro* water stress over the elements of growth and development

Variety	Inducer medium of <i>in vitro</i> water stress	The concentration of water stress simulator (%)	The average number of leaves	The average number of internodes	The average height of plantlet (cm)	The average length of root (cm)	The average weight of fresh plantlet (mg)	The average weight of fresh root (mg)		
Ruxandra	Control medium (MS)	-	8.25 CDEFG	6.75 A	11.375 B	7.875 BCDE	244,700 A	163,625 A		
	MS medium+PEG	0.5	7.00 FGHJ	5.25 CDEFG	9.075 CD	7.875 BCDE	165,675 F	76,350 EF		
		1	7.50 GHIJKL	5.00 DEFGH	8.625 CDE	7.125 CDEFGH	153,200 G	83,300 D		
	MS medium +sorbitol	1.5	7.75 EFGHI	5.75 ABCDE	8.25 CDEF	PQRST	100,125 KL	41,950 NO		
		2	6.75 IJKL	3.75 IJ	5.375 IJ	8.000 BCD	85,225 OP	52,650 HIJ		
	Sarmis	MS medium +sorbitol	0.5	8.00 DEFGH	5.50 BCDEF	7.925 EFG	6.000 HLM	187,775 D	123,625 B	
		Control medium (MS)	1	8.75 CDE	5.50 BCDEF	6.800 HIJ	8.125 BC	113,95 I	100,300 C	
			2	6.25 KL	5.0 DEFGH	5.000 KLM	8.000 BCD	81,100 M	79,600 DE	
		MS medium + PEG	-	8.00 DEFGH	6.25 ABC	13.625 A	TU	23,525	33,800	PQ
			0.5	7.25 GHIJK	5.00 DEFGH	9.125 C	4.625 NOPQRS	218,650 B	72,525 F	
1			7.25 GHIJK	4.50 FGH	4.850 MNOPQR	4.900 OPQRST	111,775 I	53,575 HI		
1.5	6.50 JKL		4.25 GHIJ	4.625 NOPQR	4.000 S	42,100 ST	47,225 KLM			
Gared	Control medium (MS)	0.5	8.00 DEFGH	4.75 EFGHI	8.750 CDE	5.625 JKLMNO	171,700 E	96,975 C		
	MS medium +sorbitol	1	8.25 CDEFGH	5.25 CDEFG	8.000 DEFG	5.675 JKLMNO	105,500 J	65,400 G		
		1.5	7.75 EFGHI	4.50 FGH	7.000 GHI	6.175 GHIJKLM	105,650 J	65,700 G		
	Control medium (MS)	2	6.00 L	3.25 J	3.300 STU	3.125 T	32,875 V	23,100 ST		
		-	7.75 EFGHI	5.00 DEFGH	7.125 GHI	7.625 BCDE	153,075 G	71,450 F		
		0.5	7.50 FGHJ	4.75 EFGHI	5.875 JKLM	5.300 KLMNOPQ	106,075 J	41,725 NO		
		1	7.00 GHIJKL	4.25 GHIJ	4.925 MNOPQR	4.900 OPQRS	50,025 R	37,525 OP		
	Marvis	MS medium + PEG	1.5	7.50 FGHJ	5.00 DEFGH	4.875 MNOPQR	7.125 CDEFGH	115,95 I	33,000 PQ	
		MS medium +sorbitol	2	7.50 FGHJ	4.00 HIJ	4.375 FGH	6.000 HIJKLM	48,000 R	24,100 S	
			1	7.00 GHIJKL	4.00 HIJ	5.625 KLMN	8.375 B	124,850 H	50,850 IJK	
Control medium (MS)		1.5	7.75 EFGHI	4.25 GHIJ	5.375 KLMNO	10,750 A	73,850 N	81,625 D		
		2	10.25 AB	5.75 ABCDE	4.000 KLMNOPQRST	7.375 BCDEF	95,250 L	66,350 G		
		-	11.00 A	6.50 AB	10.375 B	7.250 BCDEFG	247,900 A	84,125 D		
	0.5	9.00 CD	6.00 ABCD	7.375 FGH	7.625 BCDEF	153,875 G	64,100 G			
Rustic	MS+PEG	1	8.75 CDE	4.50 FGH	4.950 KLM	4.000 RSTU	67,325 O	24,175 S		
	MS+sorbitol	1.5	9.25 BC	4.75 EFGHI	4.875 MNOPQR	4.625 NOPQRS	46,775 RS	13,125 V		
		2	10.50 A	5.25 CDEFG	2.975 TU	3.700 STU	21,600 W	5,925 W		
	Control medium (MS)	0.5	6.50 KL	4.75 EFGHI	7.250 FGH	6.250 FGHJKL	124,950 H	82,375 D		
		1.5	11.25 A	6.50 AB	3.825 JKLMN	5.750 FGHJKL	104,350 JK	32,950 C		
		2	7.75 EFGHI	4.50 FGH	3.950 KLMNOPQRST	6.250 FGHJKL	57,600 Q	16,950 UV		
-		9.00 CD	5.50 BCDEF	11.00 B	7.750 BCDE	173,350 E	56,250 H			
Rustic	MS medium + PEG	0.5	8.50 CDEF	5.50 BCDEF	7.175 FGH	LMNOPQR	183,600 D	45,425 LMN		
	MS medium +sorbitol	1	8.25 CDEFG	4.50 FGH	6.375 HIJK	4.500 OPQRS	103,775 JK	29,625 OR		
		1.5	8.25 CDEFG	4.75 EFGHI	4.950 MNOPQ	4.125 QRST	57,550 K	18,675 TU		
	Control medium (MS)	2	6.25 KL	3.25 J	2.750 U	3.075 T	64,800 OP	6,275 W		
		0.5	8.50 CDEF	5.25 CDEFG	7.125 GHI	6.675 DEFGHI	156,150 G	42,550 MN		
		1	8.75 CDE	5.75 ABCDE	6.125 IJKL	5.625 JKLM	195,825 C	48,475 JKL		
1.5		7.75 EFGHI	5.00 DEFGH	5.250 LMNOP	4.750 KLMNOP	102,625 JK	27,200 RS			
2	8.75 CDE	5.00 DEFGH	4.175 PQRS	4.750 NOPQRS	45,900 RS	33,825 PQ				

LSD=4,994 mg

LSD=5,047 mg

LSD=1,186 cm

LSD=1,116 cm

LSD=1,105

LSD=1,246

Means that are in the same column followed by the same letters are not significant, according to Duncan test ($p \leq 0.05$).

The medium that had in composition PEG which is a more appropriate simulator of water stress *in vitro* determined for varieties studied lower values compared with sorbitol.




Figure 3 Effects of water stress stimulators (PEG and sorbitol) on average height of plantlet and root length

The smallest influence of water stress on plantlets height (figure 3) was observed for Ruxandra variety. By using PEG in culture medium, plantlets reached 7.83 cm. For Sarmis variety was observed a very high ratio between height plantlets developed on control medium and medium with drought simulators (MS medium: MS+PEG medium the ratio is 2.55 and MS medium: MS+sorbitol medium ratio is 2.02), so we can conclude low tolerance drought.

A good tolerance to water stress for root length, presented Gared and Ruxandra varieties when was applied sorbitol on growth medium. The root plantlets reached an average length of 8.13 and 7.22 cm. On addition of PEG, it appears that the same varieties Ruxandra and Gared, shows tolerance to water stress, but being a stronger agent, the levels were lower.




Figure 4 Effects of water stress stimulators (PEG and sorbitol) on average weight of fresh plantlet and root

Regarding the average weight of plantlet (figure 4) the highest values were observed for Ruxandra variety (121.06 cm) by applying PEG on growth medium and Rustic variety (125.13 cm), by using sorbitol. This varieties presented tolerance to drought.

CONCLUSIONS

Experimental results showed that the sensitivity to drought was not uniform for the analyzed varieties.

Medium in which was added PEG in different concentrations significantly reduced the average weight of fresh plantlet and fresh root compared with the control medium and sorbitol and significantly reduced the mean number of internodes, the average height of the plantlet, the average root length. This osmotic agent (PEG) is recommended for simulation *in vitro* of drought and to identify genotypes tolerant to drought.

In the process of leaves formation Marvis variety is highlighted, showing both for using of control medium, the greater average number of leaves (11) and on nutrient media which is applied to PEG (9.38 leaves) and sorbitol (8.31).

Regarding the average number of internodes, this decreases with the addition of PEG and sorbitol in culture medium.

The influence of variety and concentrations of osmotic agents marks Marvis variety, who under the influence of concentration of 1.5% of sorbitol has the highest value of the average number of leaves (11.25).

The smallest influence of water stress on plantlets height was observed for variety Ruxandra.

Gared and Ruxandra varieties showed good tolerance to water stress on root length, by applying sorbitol in nutritive medium, the root of plantlets reaching an average length of 8.13 and 7.22 cm.

By adding PEG (1.5%) Gared variety is characterized by achieving high values of the average weight of fresh plantlet (115.950 mg).

For average root weight is distinguished Gared variety that by using sorbitol 1% records a value of 81.625 mg (higher than obtained for control medium: 71.45 mg).

For the parameters analyzed, Marvis Gared, Ruxandra varieties, showed the best *in vitro* tolerance to water stress.

REFERENCES

Morar Gavrilă, 1999 - *Cultura cartofului*, Editura Risoprint, Cluj-Napoca.

Murashige T., Skoog F., 1962 - A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 15, p. 473-497.

Obidiegwu Jude E., Bryan Glenn J., Jones Hamlyn G., Prashar Ankush, 2015 - *Coping with drought: stress and adaptive responses in potato and perspectives for improvement*. *Frontiers in Plant Science*; Vol. 6, Art. 542, p. 1-23.

Petcu Elena, 2008: Impactul schimbărilor climatice asupra plantelor: seceta, Editura Dominor.

Popovici A., Negru P., Șișcanu Gh, 2013 - *Influența stresului hidric și hipotermic asupra metabolismului compușilor fosforici la plantele*

viticole, *Fiziologia și Biochimia Plantelor*, Buletinul AȘM. Științele vieții. Nr. 2 (320).

Săulescu, N.A, Săulescu, N.N., 1967 - *Cîmpul de experiență, Întreprinderea poligrafică Sibiu*.

Tardieu F., 1994 - *Growth and functioning of roots and of root systems subject to compaction towards a system with multiple signalling*. *Soil and Tillage Research* 30, p. 217-43.

Usman Kris Joko Suharjo, 2012 - *The effect of Polyethylene Glycol (PEG) 8000 on the growth of seven potato genotypes and their tuber production in vitro*. *Proc Soc Indon Biodiv Int*, Conf. vol. 1, p. 38-42.

PRINT ISSN: 1454-7414
ELECTRONIC ISSN: 2069-6727
CD - ROM 2285-8148

UNIVERSITATEA DE ȘTIINȚE AGRICOLE
ȘI MEDICINĂ VETERINARĂ
“ION IONESCU DE LA BRAD” IAȘI

LUCRĂRI ȘTIINȚIFICE

SERIA AGRONOMIE

VOL. 59, NR. 1

EDITURA “ION IONESCU DE LA BRAD” IAȘI



2016

COORDONATORII REVISTEI „LUCRĂRI ȘTIINȚIFICE”

Redactor responsabil: Prof. dr. Vasile VÎNTU – USAMV Iași

Redactor adjunct: Prof. dr. Liviu MIRON – USAMV Iași

Membri:

- Prof. dr. Costel SAMUIL – USAMV Iași
- Prof. dr. Lucia DRAGHIA – USAMV Iași
- Prof. dr. Paul BOIȘTEANU – USAMV Iași
- Prof. dr. Gheorghe SAVUȚA – USAMV Iași

COLEGIUL DE REDACȚIE AL SERIEI „AGRONOMIE”

Redactor șef: Prof. dr. Costel SAMUIL – USAMV Iași

Redactor adjunct: Șef lucrări dr. Florin LIPȘA – USAMV Iași

Redactori:

- Prof. dr. Wolfgang FRIEDT – Justus-Liebig-Universität Giessen, Germany
- Prof. dr. Bernd HONERMEIER – Justus-Liebig-Universität Giessen, Germany
- Prof. dr. Gerard JITĂREANU – USAMV Iași, România
- Prof. dr. Christos TSADILAS – Institute of Soil Mapping and Classification Larissa, Greece
- Prof. dr. Ioan ȚENU – USAMV Iași, România
- Prof. dr. Ioan ROTAR – USAMV Cluj Napoca, România
- Prof. dr. Roxana VIDICAN – USAMV Cluj Napoca, România
- Prof. dr. Florin IMBREA - USAMV Timișoara, România
- Prof. dr. Karoly BODNAR – Universitatea din Szeged, Ungaria
- Prof. dr. Vito Nicola SAVINO – Universitatea de Studii din Bari, Italia
- Conf. dr. Ion BĂCEAN – Universitatea Agrară de Stat din Chișinău, Republica Moldova
- Prof. dr. Jan MOUDRY Jr. – University of South Bohemia in České Budějovice, Faculty of Agriculture, Czech Republic
- Prof. dr. Petr KONVALINA - University of South Bohemia in České Budějovice, Faculty of Agriculture, Czech Republic
- Prof. dr. Gheorghe CIMPOIEȘ – Universitatea Agrară de Stat din Chișinău, Republica Moldova
- Prof. dr. Boris BOINCEAN – Universitatea Alecu Russo din Bălți, Republica Moldova
- Prof. dr. Andrei BALINSKY – Universitatea Alecu Russo din Bălți, Republica Moldova
- Prof. dr. Adriano CIANI – Perugia University
- Prof. dr. Carmen del CAMPILLO GARCIA – University of Cordoba, Campus of Rabanales, Spania
- Dr. Virginijus FEIZA – Lithuanian Institute of Agriculture, Lithuania

REFERENȚI ȘTIINȚIFICI:

- Prof. dr. Ioan AVARVAREI – USAMV Iași
- Prof. dr. Mihail AXINTE – USAMV Iași
- Prof. dr. Daniel BUCUR – USAMV Iași
- Prof. dr. Aurel CHIRAN – USAMV Iași
- Prof. dr. Viorica IACOB – USAMV Iași
- Prof. dr. Gerard JITĂREANU – USAMV Iași
- Prof. dr. Paul SAVU – USAMV Iași
- Prof. dr. Teodor ROBU – USAMV Iași
- Prof. dr. Carmenica Doina JITĂREANU – USAMV Iași
- Prof. dr. Gheorghe ȚĂRDEA – USAMV Iași
- Prof. dr. Eugen ULEA – USAMV Iași
- Prof. dr. Vasile VÎNTU – USAMV Iași
- Prof. dr. Teodor IACOB – USAMV Iași
- Prof. dr. Costel SAMUIL – USAMV Iași
- Prof. dr. Costică AILINCĂI – USAMV Iași
- Prof. dr. Stejărel BREZULEANU – USAMV Iași
- Prof. dr. Culiță SÎRBU – USAMV Iași
- Prof. dr. Mihai STANCIU – USAMV Iași
- Prof. dr. Elena GÎNDU – USAMV Iași
- Prof. dr. Gavril ȘTEFAN – USAMV Iași
- Conf. Dr. Feodor FILIPOV – USAMV Iași
- Șef lucrări dr. Florin LIPȘA – USAMV Iași

ISSN 1454-7414

Editat cu sprijinul Ministerului Educației Naționale și Cercetării Științifice

© Editura „Ion Ionescu de la Brad” Iași

Lucrări Științifice - vol. 59(1) 2016, seria Agronomie

COORDINATORS OF JOURNAL “LUCRĂRI ȘTIINȚIFICE”

Manager Editor: PhD, Prof. Vasile VÎNTU – USAMV Iași

Assistant Editor: PhD, Prof. Liviu MIRON – USAMV Iași

Members:

- PhD, Prof. Costel SAMUIL – USAMV Iași
- PhD, Prof. Lucia DRAGHIA – USAMV Iași
- PhD, Prof. Paul BOIȘTEANU – USAMV Iași
- PhD, Prof. Gheorghe SAVUȚA – USAMV Iași

EDITORIAL BOARD OF “AGRONOMIE”

Editor-in-chief: PhD, Prof. Costel SAMUIL – USAMV Iași

Assistant Editor: PhD, Lecturer Florin LIPȘA – USAMV Iași

Editors:

- PhD, Prof. Wolfgang FRIEDT – Justus-Liebig-Universität Giessen, Germany
- PhD, Prof. Prof. dr. Bernd HONERMEIER – Justus-Liebig-Universität Giessen, Germany
- PhD, Prof. Gerard JITĂREANU – USAMV Iași, Romania
- PhD, Prof. Christos TSADILAS – Institute of Soil Mapping and Classification Larissa, Greece
- PhD, Prof. Ioan ȚENU – USAMV Iași, România
- PhD, Prof. Ioan ROTAR – USAMV Cluj Napoca, Romania
- PhD, Prof. Roxana VIDICAN – USAMV Cluj Napoca, Romania
- PhD, Prof. Florin IMBREA – USAMVB Timișoara, Romania
- PhD, Prof. Karoly BODNAR – Universitatea din Szeged, Ungaria
- PhD, Prof. Vito Nicola SAVINO – Universitatea de Studii din Bari, Italia
- PhD, Assoc. Prof. Ion BĂCEAN – State Agricultural University from Chișinău, Republic of Moldova
- PhD, Prof. Jan MOUDRY Jr. – University of South Bohemia in České Budějovice, Czech Republic
- PhD, Prof. Petr KONVALINA – University of South Bohemia in České Budějovice, Czech Republic
- PhD, Prof. Gheorghe CIMPOIEȘ – Universitatea Agrară de Stat din Chișinău, Republic of Moldova
- PhD, Prof. Boris BOINCEAN – Universitatea Alecu Russo din Bălți, Republic of Moldova
- PhD, Prof. Andrei BALINSKY – Universitatea Alecu Russo din Bălți, Republic of Moldova
- PhD, Prof. Adriano CIANI – Perugia University, Italy
- PhD, Prof. Carmen del CAMPILLO GARCIA – University of Cordoba, Campus of Rabanales, Spania
- PhD. Virginijus FEIZA – Lithuanian Institute of Agriculture, Lithuania

SCIENTIFIC REVIEWERS:

- PhD, Prof. Ioan AVARVAREI – USAMV Iași
- PhD, Prof. Mihail AXINTE – USAMV Iași
- PhD, Prof. Daniel BUCUR – USAMV Iași
- PhD, Prof. Aurel CHIRAN – USAMV Iași
- PhD, Prof. Viorica IACOB – USAMV Iași
- PhD, Prof. Gerard JITĂREANU – USAMV Iași
- PhD, Prof. Paul SAVU – USAMV Iași
- PhD, Prof. Teodor ROBU – USAMV Iași
- PhD, Prof. Carmenica Doina JITĂREANU – USAMV Iași
- PhD, Prof. Gheorghe ȚĂRDEA – USAMV Iași
- PhD, Prof. Eugen ULEA – USAMV Iași
- PhD, Prof. Vasile VÎNTU – USAMV Iași
- PhD, Prof. Teodor IACOB – USAMV Iași
- PhD, Prof. Costel SAMUIL – USAMV Iași
- PhD, Prof. Costică AILINCĂI – USAMV Iași
- PhD, Prof. Stejărel BREZULEANU – USAMV Iași
- PhD, Prof. Culiță SÎRBU – USAMV Iași
- PhD, Prof. Mihai STANCIU – USAMV Iași
- PhD, Prof. Elena GÎNDU – USAMV Iași
- PhD, Prof. Gavril ȘTEFAN – USAMV Iași
- PhD, Assoc. Prof. Feodor FILIPOV – USAMV Iași
- PhD, Lecturer Florin LIPȘA – USAMV Iași

ISSN 1454-7414

Editat cu sprijinul Ministerului Educației Naționale și Cercetării Științifice

© Editura „Ion Ionescu de la Brad” Iași

Lucrări Științifice - vol. 59(1) 2016, seria Agronomie

CONTENTS

Onder UYSAL, Kamil EKİNCİ THE EFFECTS OF NITROGEN DEFICIENCY ON THE PRODUCTION OF POLYHYDROXYBUTYRATE (PHB) USING <i>SPIRULINA PLATENSIS</i>	11
Onder UYSAL, F. Ozge UYSAL, Kamil EKİNCİ DETERMINATION OF FERTILIZING CHARACTERISTICS OF THREE DIFFERENT MICROALGAE CULTIVATED IN RACEWAYS IN GREENHOUSE CONDITIONS	15
Roxana VIDICAN, Vlad STOIAN, Ioan ROTAR, Florin PĂCURAR, Susana SFECHIȘ OSCILLATION INDUCED BY AGRICULTURAL INPUTS IN MICROBIAL COMMUNITIES FORMED IN SOYBEAN RHIZOSPHERE	19
Carmen Gabriela MANOLE, Maria PARASCHIV, Laura MARIN, Mugurași CONSTANTIN, Liliana BĂDULESCU RAGWEED AND BIRTHWORT – ANTIMICROBIAL EVALUATION	25
Aurica Breica BOROZAN, Diana RABA EVALUATION OF BACTERIAL COMMUNITY AND OF ACTINOMYCETES COMPOSITION FROM THE <i>ALLIUM URSINUM</i> L SPECIES RHIZOSPHERE	31
Sándor RÓZSA, Dănuț-Nicolae MĂNIUȚIU, Rodica SIMA, Tincuța-Marta GOCAN, Ileana ANDREICA RESEARCH ON THE INFLUENCE OF HYBRID, CULTURE SUBSTRATE AND METHOD OF DISINFECTION ON OYSTER MUSHROOMS - <i>PLEUROTUS</i> SPP.	35
Emanuel VAMANU THE EFFECT OF A CONSORTIUM PROBIOTIC ADMINISTRATION ON A MICROBIOTA MICROBIOLOGICAL STABILITY	39
Carmen Liliana BĂDĂRĂU, Sorin Claudian CHIRU, Elena RAKOSY TICAN, Adriana AURORI, Gheorghe OLTEANU, Adrian GHINEA, Andreea NISTOR, Mihaela CIOLOCA PRELIMINARY STUDIES REGARDING THE INCIDENCE OF POTATO VIRUS Y IN SEED POTATOES IN ROMANIA (FOR SEVERAL CULTIVARS)	43
Maria DUCA, Steliana CLAPCO, Angela PORT ANALYSIS OF THE SOIL PARAMETERS IN THE CONTEXT OF SUNFLOWER INFECTION BY <i>Orobanche cumana</i> WALLR.	49
Liviu GRUIA, Cristinel Relu ZALĂ, Stelica CRISTEA RESEARCHES ON THE EFFICIENCY OF SOME FUNGICIDES IN MUSTARD ALTERNARIA BLACK SPOT CONTROL	53

Andreea-Mihaela BĂLĂU, Florin Daniel LIPȘA, Eugen ULEA EVALUATION OF SOME WHEAT VARIETIES BEHAVIOR TO INFECTION WITH TILLETIA CARIES (D.C.) TUL.	57
Cătălina-Ioana IACOB, Andreea-Mihaela BĂLĂU, Florin-Daniel LIPȘA, Eugen ULEA THE INFLUENCE OF PDA, CMA, TA AND V8 CULTURE MEDIA ON THE DEVELOPMENT OF PHYTOPHTHORA INFESTANS DE BARY. AND PHYTOPHTHORA PARASITICA DAST.	63
Sergiu PANUȚA, Nicolaie CROITORU, Oana Maria LĂCĂTUȘU (BODESCU) NEW MEANS OF COMBATING PHYTOPHTHORA INFESTANS AND ALTERNARIA SOLANI FUNGI IN POTATO CROP	69
Handan ÇULAL-KILIÇ, Nejla YARDIMCI MOLECULAR AND SEROLOGICAL DETECTION OF CUCUMBER MOSAIC VIRUS IN PEPPER GROWING AREAS IN ISPARTA PROVINCE, TURKEY	73
Eugen VELICHI THE DYNAMICS OF THE OCCURRENCE AND EVOLUTION OF THE ATTACK OF SOME PATHOGENIC AGENTS ON MELONS IN THE PEDOCLIMATIC CONDITIONS OF THE NORTH – EASTERN BĂRĂGAN IN 2016	77
Nela TĂLMACIU, Mihai TĂLMACIU, Monica HEREA, Ionela MOCANU RESEARCH ON THE FAUNA OF CARABIDAE (COLEOPTERA, CARABIDAE) OF SOME PLUM ORCHARDS IN EASTERN ROMANIA	81
Emil GEORGESCU, Radu GARGARITĂ, Lidia CANĂ, Luxita RASNOVEANU PRELIMINARY STUDY CONCERNING CLIMATIC CONDITIONS INFLUENCE FROM WINTER SEASON ON MAIZE LEAF WEEVIL (TANYMECUS DILATICOLLIS GYLL) ATTACK	87
Olesea COJOCARU, Rodica MELNIC EARTHWORMS (LUMBRICIDAE) - AN IMPORTANT ROLE IN SOIL FORMATION	93
Andreea TICAN, Mihaela CIOLOCA, Nicoleta CHIRU, Carmen BĂDĂRĂU BEHAVIOUR OF DIFFERENT POTATO VARIETIES BY SIMULATING IN VITRO OF HYDRIC STRESS CONDITIONS	97
Petr KONVALINA, Marek KOPECKÝ, Jaroslav BERNAS, Jan MOUDRÝ Jr., Karel SUCHÝ, Božena MACHKOVÁ WEED COMPETITIVENESS OF WINTER AND SPRING WHEAT IN ORGANIC FARMING	103
Marek KOPECKÝ, Jan MOUDRÝ Jr., Jaroslav BERNAS, Zuzana JELÍNKOVÁ, Petr KONVALINA, Radka VÁCHALOVÁ, Jan MOUDRÝ, Ladislav KOLÁŘ EFFECT OFFERTILIZATION ON THEYIELDS OF TALL WHEATGRASS HARVESTED ONCE A YEAR	109

Nina BĂRĂSCU, Maria IANOSI, Edward MUNTEAN, Marcel DUDA, Victor DONESCU THE NPK FERTILIZATION EFFECTS ON CULINARY AND TECHNOLOGICAL POTATO QUALITY	113
Radu Hermeziu, Manuela Hermeziu, Maria Stefan COSIANA – NEW ROMANIAN POTATO VARIETY	119
Susana Felicia BUTUZA BUMB, E. LUCA, Tincuța BOTA GOCAN, Sandor ROZSA THE INFLUENCES OF THE IRRIGATION, FERTILIZATION AND VARIETIES IN OBTAINING THE ONION CROP, IN MIHAI VITEAZU AREA, CLUJ COUNTY, IN 2014 ...	123
Susana Felicia BUTUZA BUMB, E. LUCA, Tincuța BOTA GOCAN, Sandor ROZSA THE INFLUENCE OF THE INTERACTION OF SOME TECHNOLOGICAL FACTORS UPON THE ONION CROP, SOWN IN 2015	127
Alexandra LEONTE, Constantin GĂUCĂ, Simona – Florina POCHIȘCANU, Alexandra – Andreea BUBURUZ, Teodor ROBU, Cătălina DRUȚU RESEARCHES REGARDING THE INFLUENCE OF „SECUIENI METHOD” ON SEED YIELDS AT MONOECIOUS HEMP (CANNABIS SATIVA L.) CULTIVATED IN THE PEDOCLIMATIC CONDITIONS OF CENTRAL MOLDAVIA.....	131
Cornelia LUPU, Simona - Florina POCHIȘCANU, Margareta NAIE, Alexandra-Andreea BUBURUZ RESEARCH REGARDING THE LONG-TERM FERTILIZATION ON THE WINTER WHEAT YIELD, AT A.R.D.S. SECUIENI	135
Simona-Florina POCHIȘCANU, Alexandra-Andreea BUBURUZ, Lorena Diana POPA, Paula Lucelia URSACHE, Alexandra LEONTE RESEARCHES REGARDING THE INFLUENCE OF SOME TECHNOLOGICAL ELEMENTS ON THE YIELD LEVELS OF SOYBEAN CROP IN THE CENTRAL OF MOLDAVIA PEDOCLIMATIC CONDITIONS PAPER TITLE.....	143
Daniela MIHALACHE, Carmen Eugenia SÎRBU, Traian Mihai CIOROIANU, Adriana Elena GRIGORE, Nicoleta MĂRIN, Ana Maria STĂNESCU, Bogdan RUJOL, Mariana IANCU STUDY ON THE EFFICIENCY AND MOBILITY OF DIFFERENT FORMS OF NITROGEN FROM FOLIAR FERTILIZERS BY USING OF ¹⁵N ISOTOPE	147
Nicoleta MĂRIN, Mihail DUMITRU, Carmen Sîrbu, Daniela MIHALACHE, Adriana GRIGORE, Ana-Maria STĂNESCU EFFICIENCY OF FOLIAR FERTILIZATION ON WHEAT PRODUCTION IN CONVENTIONAL AND CONSERVATION AGRICULTURE.....	153
Despina-Maria BORDEAN, Diana RABA, Simion ALDA, Sofia PINTILIE, Aurica Breica BOROZAN, Monica HARMANESCU, Luminita PIRVULESCU MUSHROOMS, THE MIRACLE OF NATURE, A TREAT TO HUMAN HEALTH	157
Andrei SIURIS, Vasile PLĂMĂDEALĂ, Tatiana CIOLACU RESEARCH ON AGRONOMIC AND ECONOMIC EFECTIVENESS OF WASTES FROM THE WINE INDUSTRY USED AS FERTILIZER IN THE REPUBLIC OF MOLDOVA	161

Monica HĂRMANESCU ORGANIC VS. MINERAL INPUT EFFECTS ON TOTAL PROTEIN CONTENT IN SUMMER OF SPONTANEOUS FLORA OF A HILL PERMANENT GRASSLAND FROM BANAT COUNTY	165
Monica HĂRMANESCU FERTILISATION IMPACT ON TOTAL FIBRE CONTENT OF A BANAT HILL GRASSLAND SPONTANEOUS FLORA IN SPRING	171
Margareta NAIE, Elena TROTUȘ, Alexandra-Andreea BUBURUZ, Cornelia LUPU, Alexandra LEONTE RESEARCH REGARDING THE BEHAVIOUR OF SOME PERENNIAL GRASSES AND LEGUMES MIXTURES IN ORDER TO ESTABLISH TEMPORARY GRASSLAND EXPLOITED IN MEADOW REGIME	177
Mihai STAVARACHE, Costel SAMUIL, Doina TARCĂU, Vasile VÎNTU1 EVOLUTION AND RELATIONSHIP OF SOME MACRO MINERALS IN <i>MEDICAGO SATIVA</i> L. PLANTS	183
Cătălina (CIOBANU) BOUREANU, Mihai STAVARACHE, Costel SAMUIL, Vasile VÎNTU INFLUENCE OF FERTILIZATION ON FORAGE QUALITY OF THE SIMPLE MIXTURES BETWEEN <i>BROMUS INERMIS</i> LEYSS. AND <i>ONOBRYCHIS VICIFOLIA</i> SCOP.....	189
Valentina ANDRIUCA, Ion BACEAN, Daniela DUBIT, Nicolai CAZMALI, Lucia MACRII, Rodica MELNIC, Vasile BEJAN THE PARTICULARITIES OF ROOT SYSTEM DEVELOPMENT IN WINTER WHEAT GROWN ON CARBONATE CHERNOZEM	193
Tamara LEAH THE INFLUENCE OF SOIL TILLAGE ON THE SOME PHYSICAL INDICATORS OF CHERNOZEM CAMBIC FROM MOLDOVA	197
Anca-Elena CALISTRU, Irina Gabriela CARA, Denis ȚOPA, Gerard JITĂREANU ANALYZING SOIL POROSITY UNDER DIFFERENT TILLAGE SYSTEMS USING X-RAY MICROTOMOGRAPHY	201
Stejărel BREZULEANU, Carmen Olguța BREZULEANU, Carmen Luiza COSTULEANU, Mihaela LUPU, Dragoș ROBU, Marius SCARLAT STRATEGIES AND TOOLS FOR DISPOSAL OF SOCIAL DISEQUILIBRIUM FROM THE RURAL AREA THROUGH GROUPS OF LOCAL ACTION	205
Vecdi DEMIRCAN, Deniz SARICA, Duygu SERT CURRENT STATUS AND DEVELOPMENT IN BEEKEEPING SECTOR IN TURKEY AND IN THE WORLD	211
Artur GOLBAN, Valeriu DOGA VALUE CHAIN ANALYSIS OF FRESH TOMATOES – CONSTRAINTS AND OPPORTUNITIES	219

Artur GOLBAN THE ANALYSIS OF THE ECONOMIC REFORMS IN THE AGRI-FOOD SECTOR AND THEIR IMPACT ON COMPETITIVENESS INCREASING OF THE MOLDOVAN AGRICULTURAL ENTERPRISES	225
Andrea FEHER, Tabita ADAMOV, Miroslav RAICOV ANALYSES OF BUSINESS DEVELOPMENT AT THE LEVEL OF TIMIS COUNTY	229
Aurel LUP, Liliana MIRON, Indira ALIM PAYMENT SYSTEMS FOR IRRIGATION WATER IN ROMANIA	235
Florin SALA, Ciprian RUJESCU, Andrea FEHER THE EVOLUTION SCENARIO FOR THE AVERAGE WHEAT YIELD IN THE WEST OF ROMANIA	239
Gabriela IGNAT, Andreea Alexandra TIMOFTE, Florentina ACOSTĂCHIOAIE GREEN ACCOUNTING VS SUSTAINABLE DEVELOPMENT	245
George UNGUREANU, Eduard BOGHITĂ, Dan BODESCU, Dan DONOSĂ, Cătălin Răzvan VÎNTU OPTIMISING PRODUCTION STRUCTURE OF AGRICULTURAL HOLDINGS AS A RESULT OF THEIR ADAPTATION TO MARKET REQUIREMENTS	249
George UNGUREANU, Eduard BOGHITĂ, Dan BODESCU, Radu MORARU, Cătălin Răzvan VÎNTU STUDIES REGARDING THE DEVELOPMENT OF AGRICULTURAL PRODUCTION IN THE NE REGION OF ROMANIA	255
Radu-Adrian MORARU, Stejărel BREZULEANU, Andrei MELNICIUC, Carmen Olguta BREZULEANU ASPECTS CONCERNING THE HUMAN RESOURCES MANAGEMENT AT SC COMFERT SRL BACĂU	261
Radu-Adrian MORARU, George UNGUREANU, Dan BODESCU, Dan DONOSĂ MOTIVATIONS AND CHALLENGES FOR ENTREPRENEURS IN AGRITOURISM	267
Alexandru-Dragoș ROBU, Stejărel BREZULEANU, Mioara BORZA, Gabriela BOLDUREANU, Marius-Cristian TOMA, Carmen Luiza COSTULEANU AN ANALYSIS OF S.C. AGROMIXT SPINENI S.R.L. AGRICULTURAL COMPANY'S EVOLUTION AS AN EUROPEAN FUNDS BENEFICIARY	273
Valeriu MOCA, Loredana CRENGANIȘ, Oprea RADU, Cristian HUȚANU, Constantin SAVU DRAWING UP THE NUMERICAL MODEL FOR THE FIELD RELIEF BASED ON THE DETAILED TOPOGRAPHIC ELEVATIONS PERFORMED IN THE UNDERGROUND PIPE DRAINAGE SYSTEMS	277
Valeriu MOCA, Cristian HUȚANU, Oprea RADU, Mihaela CÂRDEI THE USE OF MAP PROJECTIONS IN THE TOPOGRAPHIC AND CADASTRAL WORKS OF IASI CITY	283

Cristian HUȚANU, Oprea RADU, Dan PĂDURE UPDATING OF CADASTRAL DATABASE WITH A VIEW TO THE CREATION OF VINEYARDS AND FRUIT TREES	289
Oprea RADU AGRICULTURAL LAND QUALITY IN ION NECULCE COMMUNE, IAȘI COUNTY	295
Oprea RADU, Cristian HUȚANU, Daniel CUREA, Oleg HORJAN LAND QUALITY AND PRODUCTIVITY ASSESSMENT DATABASE CONCERNING THE AGRICULTURAL LAND OF ION NECULCE COMMUNE, IAȘI COUNTY	301
Radu ROȘCA, Ciprian CHIRUȚĂ, Virgil VLAHIDIS, Radu George CIORAP PRELIMINARY TESTS REGARDING THE OPERATING CONDITIONS AND MODELING OF THE ENERGY CONSUMPTION FOR A DOMESTIC REFRIGERATOR	307
Constantin CHIRILĂ, Dan CAZACU INFLUENCE OF SLIPPING OF DRIVE BELT FROM DISTRIBUTION DEVICE AND SLIPPING OF SUPPORTING WHEEL, DRIVING WHEEL OF DISTRIBUTION DEVICES ON SOWING PRECISION, IN CASE OF A SOWING MACHINE FOR EXPERIMENTAL FIELDS	313